

New reports, phylogenetic analysis, and a key to *Lactarius* Pers. in the Greater Yellowstone Ecosystem informed by molecular data

Edward G. Barge¹, Cathy L. Cripps¹

¹ 119 Plant Biosciences Building, Plant Science and Plant Pathology Department, Montana State University, Bozeman, MT, USA 59717

Corresponding author: Edward G. Barge (ebarge9@gmail.com)

Academic editor: K. Hosaka | Received 16 June 2016 | Accepted 25 August 2016 | Published 8 September 2016

Citation: Barge EG, Cripps CL (2016) New reports, phylogenetic analysis, and a key to *Lactarius* Pers. in the Greater Yellowstone Ecosystem informed by molecular data. MycoKeys 15: 1–58. doi: 10.3897/mycokeys.15.9587

Abstract

The Greater Yellowstone Ecosystem (GYE), located in the Central Rocky Mountains of western North America, is one of the largest nearly intact temperate-zone ecosystems on Earth. Here, *Lactarius* is an important component of ectomycorrhizal communities in many habitat types, from low elevation riparian areas to high elevation conifer forests and alpine tundra. Molecular phylogenetic analyses of ITS and *RPB2* gene sequences along with detailed morphological examination confirm at least 20 *Lactarius* species, as well as three varieties, and one unresolved species group in the GYE. Eight taxa are reported from the GYE for the first time, and nearly every major ectomycorrhizal host plant in the GYE appears to have at least one *Lactarius* species associated with it. Broad intercontinental distributions are suggested for alpine *Salix* and *Betula* associates, and for certain subalpine *Picea* and aspen (*Populus* spp.) associates. Some species appear to be restricted to western North America with *Pinus*, *Pseudotsuga* or *Abies*. The distribution and/or host affinities of others is not clear due in part to ambiguous host assignment, taxonomic problems or the relative rarity with which they have been reported.

Key words

ITS, *RPB2*, Russulaceae, Rocky Mountains, fungal biodiversity, ectomycorrhizal, systematics

Introduction

The Greater Yellowstone Ecosystem (GYE), located in the Central Rocky Mountains of North America, is one of the largest nearly intact temperate-zone ecosystems on Earth. While a definitive boundary does not exist, the GYE roughly includes Yellowstone National Park at its center, Grand Teton National Park, and portions of surrounding national forests and other lands in Montana, Wyoming, and Idaho (Keiter and Boyce 1991) (Figure 1). The Rocky Mountains, which stretch from Northern Alaska to Northern New Mexico make up the defining landform upon which the ecosystem rests. The mosaic of geology and microclimates created by the Rocky Mountains form a variety of distinct habitats for plants and fungi in the GYE from low elevation grasslands and sagebrush steppe to high elevation alpine tundra (Marston and Anderson 1991, Knight 1994).

The genus *Lactarius* Pers. (Russulales, Russulaceae) is an important component of ectomycorrhizal fungal communities throughout the Rocky Mountains, where it associates with most major ectomycorrhizal plant families. *Lactarius* as originally defined, includes species with sporocarps that exude a milky latex when damaged, in addition to the amyloid ornamented basidiospores characteristic of the family. The original genus is now recognized as nonmonophyletic and has been split into three genera within Russulaceae: *Lactarius*, *Lactifluus* (Pers.) Roussel, and *Multifurca* Buyck & Hoffstetter (Buyck et al. 2008, 2010); although approximately 80% of the traditional species are still retained in *Lactarius* sensu stricto (Verbeken and Nuytinck 2013).

Lactarius is well-documented from the Rocky Mountains and western North America (Hesler and Smith 1979, Laursen and Ammirati 1982, Methven 1997, Bessette et al. 2009, Geml et al. 2009). For alpine areas of the Rocky Mountains, including material from the GYE, Cripps and Barge (2013) and Barge et al. (2016) delineated six to seven species, most with broad, intercontinental distributions using molecular and morphological techniques. Other previous reports of *Lactarius* from the GYE specifically include McKnight (1982), Moser and McKnight (1987), Cripps and Miller (1993), Cripps (2001, 2003, 2011), Mohatt et al. (2008), Cripps and Horak (2008), Cripps and Antibus (2011), Cripps and Eddington (2005, 2012), and Cripps et al. (2016). While the above-cited works provide crucial information on *Lactarius* occurring in western North America and the Rockies, we felt it would be beneficial to provide an updated summary of the genus in the Greater Yellowstone Ecosystem as a whole, report new records, add phylogenetic information, and bring to light groups which may require further taxonomic clarification.

For this study, collections of *Lactarius* from the GYE are evaluated through detailed morphological study and phylogenetic analyses of sequences from two genetic loci: nuclear rDNA ITS1–5.8S–ITS2 (ITS region) and the region between conserved domains six and seven of the second largest subunit of the RNA polymerase II (*RPB2*) gene. Data were compared to type specimens and specimens from type localities where available, and to related taxa represented by sequences in databases (GenBank, UNITE). Information previously generated in Barge et al. (2016) was also included and forms the backbone of this study. A key to species, morphological descriptions,




Figure 1. Map showing the location of the Greater Yellowstone Ecosystem (GYE). The GYE is located in the Central Rocky Mountains of North America and includes over 20 mountain ranges, with Yellowstone National Park at its center, Grand Teton National Park, and portions of surrounding national forests and other lands in Montana, Wyoming, and Idaho. The GYE is outlined by the black box, and the Rocky Mountains by the dotted line.

insight into ecology and distribution, as well as species photographs are provided. The key should also be relevant elsewhere in the Rocky Mountains, however, less so where host plants not present in the GYE occur, such as Northwestern Montana, and Northern Idaho. Genus-specific guides covering western North American species include Hesler and Smith (1979), Methven (1997), Bessette et al. (2009), and the Pacific Northwest Key Council's Key to *Lactarius* species in the Pacific Northwest (<http://www.svims.ca/council/Lactar.htm>).

Methods

Study sites

This study was performed in the GYE (Figure 1), with a focus on northern portions of the ecosystem. Major ectomycorrhizal host plants targeted include *Populus* spp. (cottonwoods) and *Alnus incana* (L.) Moench (gray alder) in riparian areas; *Pinus flexilis* James (limber pine), *Pseudotsuga menziesii* (Mirb.) Franco (Douglas fir), *Cercocarpus ledifolius* Nutt. (curl-leaf mountain mahogany) and *Populus tremuloides* Michx. (quaking aspen) in low to mid-elevation montane areas; *Pinus contorta* Dougl. ex Loud.

(lodgepole pine), *Pinus albicaulis* Engelm. (whitebark pine), *Picea engelmannii* Parry ex Engelm. (Engelmann spruce), *Abies lasiocarpa* (Hook.) Nutt. (subalpine fir), *Betula glandulosa* Michx. (dwarf birch), and *Arctostaphylos uva-ursi* (L.) Spreng. (kinnikinnick) in mid to high-elevation montane areas; and dwarf willows (*Salix arctica* Pall., *S. reticulata* L.), shrubby willows (*S. glauca* L., *S. planifolia* Pursh), *Dryas octopetala* L. (mountain avens), *Betula glandulosa*, and *Arctostaphylos uva-ursi* at or above treeline. *Pinus ponderosa* Dougl. ex Laws., which occurs sporadically along the northeastern edge of the ecosystem, was not surveyed. The locations of specimens are included in the Taxonomy section under Specimens examined and locations of specimens used in the phylogenetic analysis are shown in Table 1.

Taxon sampling and processing

Sporocarps were collected from late May to late September 1992–2015. All collections were described in detail when fresh, and select collections were photographed. Ectomycorrhizal host plants near sporocarps were noted for each collection. Sporocarps were dried on an electric, warm-air dryer and deposited in the Montana State University herbarium (MONT). For comparison, additional specimens were obtained from numerous herbaria. For Rocky Mountain material, most herbarium specimens studied originated from the Montana State University Herbarium (MONT), the Denver Botanic Gardens' Sam Mitchel Herbarium of Fungi, Denver, Colorado (DBG), and the University of Michigan fungal herbarium, Ann Arbor, Michigan (MICH). See Barge et al. (2016), for a complete list of herbaria.

Morphological descriptions

Macromorphological measurements and descriptions were made from fresh material. Micromorphological descriptions and measurements were made from dry material reconstituted in ethanol and 2.5% KOH. In Melzer's reagent, length and width were measured at 1000× magnification for a random sample of 25 basidiospores per collection, excluding ornamentation and the hilar appendix. The length to width ratio (Q) was calculated for each basidiospore. Length and width were measured at 400× magnification for a random sample of 10 pleuromacrocyttidia and 10 cheilomacrocyttidia per collection. Morphological descriptions for alpine specimens were previously reported in Barge et al. (2016).

Phylogenetic analysis

DNA extraction, PCR amplification, and sequencing of the ITS region and partial *RPB2* gene were performed as in Barge et al. (2016) or at ALVALAB, Santander, Spain

Table 1. Voucher, locality information and GenBank or UNITE (UDB) accession numbers for DNA sequences used in the phylogenetic analysis. Herbarium acronyms follow Thiers <http://sweetgum.nybg.org/ih/> (continuously updated). Specimens from the Greater Yellowstone Ecosystem (GYE) are in bold type. Newly generated accessions are in bold type.

Taxon	Voucher	Location	ITS	RPB2
<i>L. akahatsu</i>	JN2004-141 (GENT)	Thailand	KF133269	KF133333
<i>L. albocarneus</i>	AV98-080 (GENT)	France	KF241545	NA
<i>L. alnicola</i>	EB0064-14 (MONT)	Gallatin Range, MT, U.S.A.	KX394276	NA
<i>L. alpinus</i> v. <i>alpinus</i>	FNL GNP-125 (DAOM)	Newfoundland, Canada	KX094937	NA
<i>L. alpinus</i> v. <i>mitis</i>	EB161-15 (MONT)	Red Lodge, MT, U.S.A.	KX394277	NA
<i>L. alpinus</i> v. <i>mitis</i>	DBG-F-021979 (DBG)	Pike National Forest, CO, U.S.A.	KX394278	KX394303
<i>L. aspideoides</i>	RL Shaffer 6957 (MICH)	Emmet Co., MI, U.S.A.	KR090893	NA
<i>L. aspideus</i>	JV24534 (TURA)	Varsinais-Suomi, Finland	KR090894	KR090972
<i>L. atroviridis</i>	AV05-306 (GENT)	U.S.A.	KF133270	KF133334
<i>L. aurantiacus</i>	JV94-422 (C)	Uppland, Sweden	KR090896	KR090974
<i>L. cf. aurantiacus</i>	JV15112F (TURA)	Nordland, Norway	KR090895	KR090973
<i>L. auriolla</i>	RW1601 (GENT)	Sweden	KF133257	KF133321
<i>L. barrowsii</i>	EB015-15 (MONT)	Story Hill, Bozeman, MT, U.S.A.	KX394279	NA
<i>L. barrowsii</i>	N Gray 91878 (RMS)	Washington, U.S.A.	EF685047	NA
<i>L. barrowsii</i>	States J AEF 1271 (MICH)	Arizona, U.S.A.	EF685046	NA
<i>L. badiosanguineus</i>	EB200-13 (MONT)	Gallatin Range, MT, U.S.A.	KX394280	NA
<i>L. badiosanguineus</i>	EB0069-14 (MONT)	Tobacco Root Mtns., MT, U.S.A.	KX394281	NA
<i>L. badiosanguineus</i>	DBG-F-017093 (DBG)	Front Range, CO, U.S.A.	KX394282	KX394304
<i>L. badiosanguineus</i>	AV04-235 (GENT)	France	KF432983	NA
<i>L. badiosanguineus</i>	AV10-044 (GENT)	Norway	KR025578	KR025655
<i>L. brunneohepaticus</i>	PAM08090315 (LIP)	Corse-du-Sud, France	HQ714726	HQ714858
<i>L. brunneoviolaceus</i>	Verbeken 04-220 (GENT)	France	KJ742392	NA
<i>L. aff. brunneoviolaceus</i>	CLC3098 (MONT)	Beartooth Plateau, MT, U.S.A.	KX394283	NA
<i>L. aff. brunneoviolaceus</i>	CLC2133 (MONT)	Finse, Norway	KR090899	KR090977
<i>L. caespitosus</i>	EB102-13 (MONT)	Gallatin Range, MT, U.S.A.	KX394284	KX394305
<i>L. caespitosus</i>	EB121-13 (MONT)	Madison Range, MT, U.S.A.	KX394285	KX394306
<i>L. camphoratus</i>	UE04.09.2004-5 (UPS)	Sweden	DQ422009	DQ421933
<i>L. chrysorrheus</i>	UE04.10.2002-8 (UPS)	Italy	KF133261	KF133325
<i>L. citriolens</i>	UE20.09.2004-03 (UPS)	Sweden	DQ422003	DQ421931
<i>L. controversus</i>	EB110-15 (MONT)	Beartooth Mtns, MT, U.S.A.	KX394286	NA
<i>L. controversus</i>	UP508	Sweden	DQ658881	NA
<i>L. controversus</i>	AV00-117 (GENT)	Italy	KF241544	NA
<i>L. cuspidaurantiacus</i>	LM4823 (XAL)	Mexico	KF891367	KF891374
<i>L. cuspidaurantiacus</i>	LM4908 (XAL)	Mexico	KF891366	KF891373
<i>L. cyathuliformis</i>	PAM08100409 (LIP)	Orne, France	HQ714738	HQ714869
<i>L. dryadophilus</i>	EL57-10 (GB)	Latnjavagge, Sweden	KR090902	KR090980
<i>L. deliciosus</i>	JN2001-046 (GENT)	Slovakia	KF133272	KF133337

Taxon	Voucher	Location	ITS	RPB2
<i>“L. deliciosus v. deterrimus”</i>	JR Herr 650 (RMS)	Wyoming, U.S.A.	EF685056	NA
<i>L. deliciosus v. olivaceosordidus</i>	J Ammirati 10762 (RMS)	Washington, U.S.A.	EF685059	NA
“L. deliciosus” group	EB0063-14 (MONT)	Gallatin Range, MT, U.S.A.	KX394287	NA
“L. deliciosus” group	EB089-15 (MONT)	Madison Range, MT, U.S.A.	KX394288	NA
“L. deliciosus” group	EB107-15 (MONT)	Hellroaring Plateau, MT, U.S.A.	KX394289	NA
<i>“L. deliciosus” group</i>	EB0047 (MONT)	Sawatch Range, CO, U.S.A.	KX394290	NA
<i>L. deterrimus</i>	UE05.09.2004-04 (UPS)	Sweden	UDB000866	NA
<i>L. evosmus</i>	UP536	Sweden	DQ58882	NA
<i>L. evosmus</i>	UE27.09.2002-1 (UPS)	France	UDB000860	NA
<i>L. flavopalustris</i>	JV23334 (TURA)	Koillismaa, Finland	KR090904	KR090982
<i>L. flexuosus</i>	UE06.09.2002-1 (UPS)	Sweden	DQ421992	DQ421925
<i>L. fulvissimus</i>	JN2012-025 (GENT)	Germany	KR025576	KR025660
L. glyciosmus	TWO269 (MONT)	Beartooth Plateau, MT, U.S.A.	KR090905	KR090983
<i>L. glyciosmus</i>	EB133 (MONT)	San Juan Range, CO, U.S.A.	KR090909	KR090986
<i>L. glyciosmus</i>	M Moser 19810148 (IB)	Femsjö, Sweden	KR090910	NA
<i>L. helvus</i>	UE08.09.2004-1 (UPS)	Sweden	KF133263	KF133327
<i>L. hepaticus</i>	JN02-049 (GENT)	Belgium	KF432980	KR025674
<i>L. hysginoides</i>	JV28432 (TURA)	Koillismaa, Finland	KR090914	KR090988
<i>L. intermedius</i>	f324 (AQUI)	Italy	UDB000368	NA
<i>L. kauffmanii</i>	F28471 (UBC)	British Columbia, Canada	KP406573	NA
<i>L. lacunarum</i>	JKLAC11092901 (GENT)	Germany	KF432982	KR025638
L. lanceolatus	CLC2358 (MONT)	Beartooth Plateau, MT, U.S.A.	KR090918	KR090992
L. lanceolatus	EB105-13 (MONT)	Beartooth Plateau, WY, U.S.A.	KR090919	KR090993
<i>L. lanceolatus</i> (Holotype)	F4239 (VPI)	Beaufort Lagoon, AK, U.S.A.	KR090915	KR090989
<i>L. lapponicus</i>	JV28335 (TURA)	Koillismaa, Finland	KX394291	NA
<i>L. leonis</i>	TU106315 (TU)	Estonia	UDB011465	NA
<i>L. lepidotus</i>	PAM08090304 (LIP)	Corse-du-Sud, France	HQ714722	HQ714854
<i>L. lignyotus</i>	UE06.09.2003-5 (UPS)	Sweden	DQ421993	DQ421926
<i>L. lilacinus</i>	EDB08101401 (LIP)	France	HQ714748	HQ714879
<i>L. lilacinus</i>	RW3774 (GENT)	Belgium	KF133275	KF133340
<i>L. luculentus v. luculentus</i>	AH Smith 79943 (MICH)	OR, U.S.A.	KR090920	NA
L. luculentus v. laetus	EB097-15 (MONT)	Crazy Mtns., MT, U.S.A.	KX394292	NA
<i>L. luculentus v. laetus</i>	DBG-F-024643 (DBG)	CO, U.S.A.	KR090922	KR090994
<i>L. luridus</i>	OB11-011 (GENT)	Belgium	KF241547	NA
<i>L. mammosus</i>	UE09.09.2004-5 (UPS)	Sweden	KF133265	NA
L. montanus	EB120-13 (MONT)	Madison Range, MT, U.S.A.	KR090925	KR090996
L. montanus	CLC3001 (MONT)	Tobacco Root Mtns., MT, U.S.A.	KR090926	KR090997
<i>L. montanus</i> (Paratype)	AH Smith 81954 (MICH)	Bonner County, ID, U.S.A.	KR090924	NA
L. nanus	EB106-13 (MONT)	Beartooth Plateau, MT, U.S.A.	KR090928	KR091000
<i>L. nanus</i>	CLC1716 (MONT)	San Juan Range, CO, U.S.A.	KR090929	KR091001
<i>L. nanus</i>	Bon 89093 (LIP)	Savoie, France	KR090935	NA
<i>L. necator</i>	AV04-231 (GENT)	France	KF133276	KF133341

Taxon	Voucher	Location	ITS	<i>RPB2</i>
<i>L. olivinus</i>	TU101411 (TU)	Estonia	UDB019698	NA
<i>L. olivinus</i>	TU106674 (TU)	Estonia	UDB015721	NA
<i>L. aff. olivinus</i>	EB0051-14 (MONT)	Silver Gate, MT, U.S.A.	KX394293	NA
<i>L. olympianus</i>	EB0070-14 (MONT)	Tobacco Root Mtns., MT, U.S.A.	KX394294	NA
<i>L. olympianus</i>	J Nuytinck 2003-032 (GENT)	U.S.A.	EF685079	NA
<i>L. pallescens</i> (Holotype)	AH Smith 81936 (MICH)	Boundary County, ID, U.S.A.	KR090938	NA
<i>L. pallidomarginatus</i> (Holotype)	EB0041 (MONT)	San Juan Range, CO, U.S.A.	KR090940	KR091010
<i>L. pseudodelicatus</i>	CLC512 (MONT)	Teton Range, ID, U.S.A.	KX394295	NA
<i>L. pseudomucidus</i>	F16301 (UBC)	British Columbia, Canada	EU486439	NA
<i>L. aff. pseudouvidus</i>	U Peintner 20040156 (IB)	Trentino, Italy	KR090942	KR091014
<i>L. pubescens</i>	EB300-15 (MONT)	Silver Bow Co., MT, U.S.A.	KX394296	NA
<i>L. pubescens</i>	CLC539 (MONT)	Cinnabar Basin, MT, U.S.A.	KX394297	NA
<i>L. pubescens</i>	CLC1611 (MONT)	Front Range, CO, U.S.A.	KX394298	NA
<i>L. pubescens</i>	UE15.09.2002-2 (UPS)	Sweden	DQ421996	DQ421929
<i>L. pubescens</i>	UP516	Sweden	DQ658884	NA
<i>L. quieticolor</i>	UE10.09.2004-1 (UPS)	Sweden	DQ422002	DQ421930
<i>L. quietus</i>	UE16.09.2004 (UPS)	Sweden	KF133264	KF133328
<i>L. repraesentaneus</i>	EB107-13 (MONT)	Beartooth Plateau, MT, U.S.A.	KR090949	KR091021
<i>L. repraesentaneus</i>	CLC1747 (MONT)	Sawatch Range, CO, U.S.A.	KR090951	KR091023
<i>L. repraesentaneus</i>	EL92-07 (GB)	Latnjavagge, Sweden	KR090953	KR091025
<i>L. romagnesii</i>	UE29.09.2006-6 (UPS)	France	DQ421989	DQ421923
<i>L. rubrilacteus</i>	EB13-040 (MONT)	Bridger Range, MT, U.S.A.	KX394299	NA
<i>L. rubrilacteus</i>	SL Miller 19-04 (RMS)	California, U.S.A.	EF685085	NA
<i>L. rufus</i>	EB125-13 (MONT)	Gallatin Range, MT, U.S.A.	KX394300	KX394307
<i>L. rufus</i>	JN2002-008 (GENT)	Norway	KF241543	NA
<i>L. rufus</i>	DG15 (AMG)	United Kingdom	UDB001601	NA
<i>L. salicis-herbaceae</i>	CLC1536 (MONT)	Sismiut, Greenland	KR090955	KR091027
<i>L. salicis-reticulatae</i>	EB0057-14 (MONT)	Beartooth Plateau WY, U.S.A.	KR090958	KR091029
<i>L. salicis-reticulatae</i>	JV15133 (TURA)	Sweden	KR090961	KR091032
<i>L. aff. salicis-reticulatae</i>	CLC1710 (MONT)	San Juan Range, CO, U.S.A.	KR090962	KR091033
<i>L. sanguifluus</i>	Hue470 (UPS)	Germany	UDB000876	NA
<i>L. scrobiculatus</i>	JN01-058 (GENT)	Slovakia	KF432968	NA
<i>L. scrobiculatus</i>	TU117089 (TU)	Latvia	UDB022822	NA
<i>L. sphagneti</i>	PL2805 (herb. P. Leonard)	United Kingdom	KF133268	KF133332
<i>L. spinosulus</i>	AT2003068 (UPS)	Sweden	KF133262	KF133326
<i>L. subdulcis</i>	JV2006-024 (GENT)	Belgium	KF133279	KF133344
<i>L. subflammeus</i> (Holotype)	AH Smith 83602 (MICH)	Tillamook County, OR, U.S.A.	KR090967	NA
<i>L. substriatus</i>	AH Smith 83693 (MICH)	Tillamook Co., OR, U.S.A.	KR090968	NA
<i>L. subviscidus</i> (Paratype)	AH Smith 83066 (MICH)	Lewis Co., WA, U.S.A.	KR090970	NA
<i>L. tabidus</i>	ED2008-026 (GENT)	Belgium	KT165309	KR025667
<i>L. thyinos</i>	A Voitek23-08-2004 (GENT)	Canada	KF133271	KF133336

Taxon	Voucher	Location	ITS	<i>RPB2</i>
<i>L. torminosus</i>	RW3183 (GENT)	Czech Republic	KF133281	KF133346
<i>L. trivialis</i>	UE27.08.2002-17a (UPS)	Sweden	DQ421991	DQ421924
<i>L. tuomikoskii</i>	TU101882 (TU)	Finland	UDB016033	NA
<i>L. aff. tuomikoskii</i>	EB052-14 (MONT)	Silvergate, MT, U.S.A.	KX394301	NA
<i>L. uvidus</i>	mh0963 (TUB)	Germany	AY606957	NA
<i>L. vietus</i>	UE11.19.2004-1 (UPS)	Sweden	KF133267	KF133331
<i>L. aff. wenquanensis</i>	LTH143 (GENT)	Thailand	EF141537	NA
<i>L. zonarioides</i>	E Bizio 2516 (MCVE)	Italy	JF908300	NA
<i>L. zonarius</i>	UE27.09.2002-4 (UPS)	France	EU278678	EU278679
<i>L. zonarius</i> v. <i>riparius</i>	CLC2933 (MONT)	Bozeman, MT, U.S.A.	KX394302	NA

(<http://www.alvalab.es/index.html>). Raw sequences were edited and contigs constructed with SeqTrace 0.8.1 (Stucky 2012).

In addition to the sequences generated in this study, select close BLAST matches to the ITS region of GYE specimens and select additional ITS and *RPB2* sequences mainly from Buyck et al. (2008), Verbeken et al. (2014), and Wisitrassameewong et al. (2016) were downloaded from GenBank (<http://www.ncbi.nlm.nih.gov/>) and UNITE (<https://unite.ut.ee/>). Many sequences generated in Barge et al. (2016) were also included.

Alignments were performed with MUSCLE (<http://www.ebi.ac.uk/Tools/msa/muscle/>) under default parameters and manually edited using PhyDE v.0.9971 (<http://www.phyde.de/index.html>). Ambiguously aligned regions of the ITS alignment were highlighted and removed with the online version of GBlocks 0.91b (Castresana 2000) under the least stringent settings. Also removed was a 268–269 base pair insertion present in the ITS1 region of several taxa (Barge et al. 2016, Rosenblad et al. 2016).

Maximum likelihood (ML) analyses were carried out using RAxML v.8 (Stamatakis 2014) on the CIPRES Science Gateway (Miller et al. 2010). Previous phylogenetic analyses by Barge et al. (2016) suggested that *Lactarius romagnesii* Bon and *L. lignyotus* Fr. of subg. *Plinthogalus* (Burl.) Hesler & A.H. Sm. form a monophyletic group distinct from subg. *Piperites* (Fr.) Kauffman and *Russularia* (Fr.) Kauffman which contain all of the ingroup species of interest in this study. Thus, *L. lignyotus* and *L. romagnesii* were selected as outgroup taxa. An ML search combined with rapid bootstrapping was run until bootstrap convergence was reached using the autoMRE option. The GTRGAMMA model was estimated separately for ITS1, 5.8S, ITS2, *RPB2* exon first, second and third codon positions, and *RPB2* intron. Single-gene and combined ML analyses were performed. Slight conflicts were detected between ITS and *RPB2* phylogenies, however, the conflicts received low support (bootstrap support <70%), thus the combined phylogeny was used. The program TreeGraph2 (Stöver and Müller 2010) was used to draw and edit the final tree.

Results

A total of 27 ITS and 5 *RPB2* sequences were generated in this study (Table 1). Based on morphological and molecular examination we report 20 *Lactarius* species, three varieties, and one putatively unresolved species group from the GYE. *Lactarius alpinus* v. *mitis* Hesler & A.H. Sm., *L. badiosanguineus* Kühner & Romagn., *L. barrowsii* Hesler & A.H. Sm., *L. aff. brunneoviolaceus* M.P. Christ., *L. luculentus* v. *laetus* Hesler & A.H. Sm., *L. aff. olivinus* Kytöv., *L. pseudodelicatus* A.H. Sm., and *L. aff. tuomikoskii* Kytöv. are reported from the GYE for the first time. In addition, the first publicly available vouchered sequences are provided for *L. alpinus* v. *mitis*, *L. lapponicus* (syn.: *L. duplicatus*), *L. pseudodelicatus*, and *L. zonarius* v. *riparius*; the first publicly available vouchered sequences from North America are provided for *L. badiosanguineus*, *L. aff. brunneoviolaceus*, *L. aff. olivinus*, and *L. aff. tuomikoskii*.

Subgenera *Piperites* and *Russularia* are well-represented in the GYE, however, species of subg. *Plinthogalus* appear to be absent, as are species of the recently segregated genus *Lactifluus*. Most major ectomycorrhizal host plants in the Greater Yellowstone Ecosystem appear to have at least one *Lactarius* species associated with them (Table 2).

The phylogenetic position of species for which molecular data was obtained is shown (Figure 2). The backbone of the phylogeny is poorly resolved, thus we refrain from

Table 2. Major ectomycorrhizal host plants in the Greater Yellowstone Ecosystem (GYE) with putatively associated *Lactarius* species. (?) denotes host plant uncertainty.

Host Plant	<i>Lactarius</i> species
<i>Abies lasiocarpa</i>	<i>L. caespitosus</i> , <i>L. luculentus</i> v. <i>laetus</i> (?)
<i>Alnus incana</i>	<i>L. alpinus</i> v. <i>mitis</i>
<i>Arctostaphylos uva-ursi</i>	" <i>L. deliciosus</i> " group
<i>Betula glandulosa</i> <i>Betula</i> spp.	<i>L. glyciosmus</i> , <i>L. pubescens</i>
<i>Cercocarpus ledifolius</i>	None detected
<i>Dryas octopetala</i>	None detected
<i>Picea engelmannii</i>	<i>L. alnicola</i> , <i>L. badiosanguineus</i> , " <i>L. deliciosus</i> " group, <i>L. luculentus</i> v. <i>laetus</i> (?), <i>L. montanus</i> , <i>L. olympianus</i> , <i>L. rufus</i> (?), <i>L. repraesentaneus</i> , <i>L. aff. olivinus</i> and <i>L. aff. tuomikoskii</i>
<i>Pinus albicaulis</i>	" <i>L. deliciosus</i> " group, <i>L. rufus</i> (?)
<i>Pinus contorta</i>	" <i>L. deliciosus</i> " group (?), <i>L. rufus</i>
<i>Pinus flexilis</i>	<i>L. barrowsii</i> , " <i>L. deliciosus</i> " group
<i>Pinus ponderosa</i>	None detected
<i>Pseudotsuga menziesii</i>	<i>L. rubrilacteus</i>
<i>Populus tremuloides</i>	<i>L. controversus</i> , <i>L. pseudodelicatus</i> , <i>L. pubescens</i>
<i>Populus</i> spp. (cottonwoods)	<i>L. zonarius</i> v. <i>riparius</i>
<i>Salix</i> spp. (alpine)	<i>L. aff. brunneoviolaceus</i> , <i>L. nanus</i> , <i>L. lanceolatus</i> , <i>L. pallidomarginatus</i> , <i>L. repraesentaneus</i> (?), <i>L. salicis-reticulatae</i>
<i>Salix</i> spp. (subalpine)	None detected




Figure 2A. Upper part of maximum likelihood phylogeny of ITS and *RPB2* molecular data. Bootstrap support values $\geq 50\%$ are shown above or below branches leading to clades. Thickened branches lead to clades receiving $\geq 70\%$ bootstrap support. Boldface type labels represent specimens from the Greater Yellowstone Ecosystem (GYE). Boldface numbers refer to the taxa treated in the Key and Taxonomy sections.

making any novel inferences regarding broad-scale infrageneric relationships. Species are organized in the Taxonomy section based on the phylogenetic results. See comments under individual species in the Taxonomy section for discussion of relationships.

Key to *Lactarius* species of the Greater Yellowstone Ecosystem

Due to dry conditions common throughout the Rocky Mountains, *Lactarius* sporocarps collected here often do not exude any noticeable latex. However, for species which have changing latex, the cut flesh often displays characteristic color changes similar to the latex or reacts with the latex to display a characteristic color, and it typi-




Figure 2B. Middle part of maximum likelihood phylogeny of ITS and *RPB2* molecular data. Bootstrap support values ≥ 50% are shown above or below branches leading to clades. Thickened branches lead to clades receiving ≥ 70% bootstrap support. Boldface type labels represent specimens from the Greater Yellowstone Ecosystem (GYE). Boldface numbers refer to the taxa treated in the Key and Taxonomy sections.




Figure 2C. Lower part of maximum likelihood phylogeny of ITS and *RPB2* molecular data. Bootstrap support values $\geq 50\%$ are shown above or below branches leading to clades. Thickened branches lead to clades receiving $\geq 70\%$ bootstrap support. Boldface type labels represent specimens from the Greater Yellowstone Ecosystem (GYE). Boldface numbers refer to the taxa treated in the Key and Taxonomy sections.

cally does so under dry conditions even when no latex is detected. Thus, if no latex is detected, examining cut surfaces of sporocarps for color change can be very useful for identifying *Lactarius* species.

- 1 Latex orange to red; cut flesh staining orange to red and eventually green... **2**
- Latex watery or whitish, unchanging or becoming violet or yellow; cut flesh unchanging or staining lilac, yellow or brownish **4**
- 2 Latex orange, often undetectable; cut flesh staining orange to \pm red and eventually green; with conifers **20. “*L. deliciosus*” group**
- Latex red, often undetectable; cut flesh quickly staining red and eventually green..... **3**
- 3 Pileus cream to pale straw yellow, lamellae dingy pinkish orange to creamy yellow-orange; with *Pinus flexilis* and possibly other *Pinus* spp..... **18. *L. barrowsii***
- Pileus light brown to yellow-orange to dingy orange; lamellae pinkish cinnamon to dull pink; with *Pseudotsuga menziesii*, and possibly *Pinus* spp..... **19. *L. rubrilacteus***

- 4 Latex unchanging or turning violet; cut flesh staining violet **5**
 – Latex unchanging or turning yellow; cut flesh unchanging or staining yellow or brownish **9**
 5 Pileus cream to yellowish **6**
 – Pileus violaceous to brownish or greyish **7**
 6 Pileus margin bearded; stipe scrobiculate; with *Picea engelmannii* and possibly *Salix glauca* in subalpine to low alpine krummholz habitats **4. *L. repraesentaneus***
 – Pileus margin glabrous; stipe non-scrobiculate; with *Salix* spp. in alpine areas **5. *L. salicis-reticulatae***
 7 Subalpine with conifers in the spruce-fir zone; pileus cuticle staining green in KOH **7. *L. montanus***
 – Alpine with *Salix*; pileus cuticle staining green in KOH or not **8**
 8 Pileus light tan to light brown; pileus cuticle unchanging in KOH **6. *L. pallidomarginatus***
 – Pileus brownish with lavender hues; pileus cuticle staining green in KOH.... **8. *L. aff. brunneoviolaceus***
 9 Pileus margin conspicuously bearded or tomentose at least when young ... **10**
 – Pileus margin glabrous or only very finely pubescent **14**
 10 Latex unchanging or turning yellow; cut flesh staining yellow; with conifers... **11**
 – Latex unchanging; cut flesh unchanging or staining brownish; with hardwoods **12**
 11 Pileus with matted fibrils especially near the margin, pale yellow-cream to yellow-tan; margin densely bearded; under *Picea engelmannii* **16. *L. aff. tuomikoskii***
 – Pileus with appressed scales of confluent hairs especially near the margin, cream to olive-buff to orange-tan; margin merely tomentose to cottony; under *Picea engelmannii*. **17. *L. aff. olivinus***
 12 Pileus azonate to faintly zonate and predominately cream to pinkish; with aspen or *Betula*..... **13. *L. pubescens***
 – Pileus zonate and predominately yellow-buff to orange-brown; with aspen or cottonwood **13**
 13 Pileus yellow-buff to pale orange-brown, lighter toward the margin; odor orange-citrusy; with aspen..... **12. *L. pseudodelicatus***
 – Pileus yellow-brown to orange brown, ± lighter toward the margin; odor sweet, fruity; with cottonwood..... **14. *L. zonarius* v. *riparius***
 14 Pileus reddish, red-brown or orange..... **15**
 – Pileus whitish, yellowish or brownish..... **20**
 15 Taste acrid **16**
 – Taste mild to slightly bitter..... **17**
 16 Pileus red-brown to orange-brown, azonate; with conifers **22. *L. rufus***
 – Pileus rich to dull orange, zonate; with conifers **11. *L. olympianus***

- 17 Pileus finely rimulose to minutely squamulose; with *Alnus*.....**1. *L. alpinus* v. *mitis***
**1. *L. alpinus* v. *mitis***
 – Pileus smooth; with conifers or *Salix*..... 18
 18 Alpine with *Salix* **24. *L. lanceolatus***
 – Subalpine with conifers.....**19**
 19 Pileus red-brown to orange-brown, often lighter toward the margin
 **21. *L. badiosanguineus***
 – Pileus bright orange to yellow-orange to orange-brown, evenly colored
**23. *L. luculentus* v. *laetus***
 20 Pileus brownish or pinkish buff; taste mild to only slightly acrid.....**21**
 – Pileus whitish, yellowish, brownish or gray-brown; taste acrid**22**
 21 Odor mild; taste mild to slightly acrid; arctic-alpine with *Salix*...**2. *L. nanus***
 – Odor of coconut; taste mild to slightly acrid; alpine to subalpine with *Betu-*
la**3. *L. glyciosmus***
 22 Pileus yellowish; flesh staining yellow; with *Picea engelmannii***15. *L. alnicola***
 – Pileus whitish, brownish or vinaceous.....**23**
 23 Pileus whitish overall; lamellae pinkish; with aspen..... **10. *L. controversus***
 – Pileus more brown to gray-brown overall; lamellae white to cream; with
Abies.....**9. *L. caespitosus***

Taxonomy

1. *Lactarius alpinus* v. *mitis* Hesler & A.H. Sm.

Figure 3

Description. Pileus 8–20 mm in diameter (to 80 mm in Hesler and Smith 1979), plano-convex to infundibuliform, \pm papillate, dry, finely rimulose to minutely squamulose, red-brown to orange-brown to pinkish cinnamon; margin finely crenulate, incurved when young, becoming straight to wavy in age. Lamellae adnate to subdecurrent, subdistant to crowded, cream to creamy tan, discoloring slightly darker where damaged. Stipe 8–23 \times 5–8 mm (30–50 \times 10–15 mm in Hesler and Smith 1979), equal to slightly clavate, smooth, dry, pale creamy yellow to creamy tan, solid, becoming hollow. Context concolorous with pileus. Latex scarce, watery, white, unchanging. Odor mild. Taste mild.

Basidiospores 7–9.5 \times 5–7.5 μ m, Q = 1.3–1.5, ellipsoid; ornamentation forming a broken to partial reticulum. Pleuromacrocytidia 60–100 \times 6–10 μ m, numerous, strongly projecting, mucronate to fusoid; apex acute to moniliform. Cheilomacrocytidia 50–80 \times 6–10 μ m, numerous, strongly projecting, mucronate to fusoid; apex acute to moniliform.

Ecology and distribution. In western and into eastern North America with *Alnus*. In the GYE only one collection has been made thus far and it was from a riparian area near *Alnus incana* and *Populus trichocarpa*, late summer. Further sampling under *Alnus* in the Rocky Mountains is necessary.



Figure 3. *Lactarius alpinus* v. *mitis*. Top collection EB161-15 under *Alnus incana* and *Populus trichocarpa*, Red Lodge, Montana, USA. Bottom collection EB516-15 under *Alnus rubra*, western Oregon, USA. Scale bars: 2 cm. Photos by E. Barge.

Specimens examined. U.S.A. MONTANA: Carbon County, Red Lodge, along Rock Creek, under *Alnus incana* and *Populus trichocarpa*, 6 Sept 2015, EB161-15 (MONT).

Discussion. This is the first report of *L. alpinus* v. *mitis* from the GYE. According to Hesler and Smith (1979), *L. alpinus* v. *mitis* has duller pileus and lamellae colors than *L. alpinus* v. *alpinus* Peck, which typically develops bright yellow to apricot colors; variety *mitis* also has longer and more conspicuous macrocystidia and a mild rather than slowly acrid taste. The collection described here likely did not reach full maturity, and the basidiomes are smaller than reported in Hesler and Smith (1979). Phylogenetic analyses including material of *L. alpinus* v. *alpinus* from near the type locality suggest that *L. alpinus* v. *mitis* and *L. alpinus* v. *alpinus* are distinct species (Figure 2A, 2C), however a more in depth analysis incorporating type material of at least *L. alpinus* v. *mitis* should be carried out before making this determination final. For an interesting discussion on *L. alpinus* see Nuytinck and Voitek (2016) and Rochet et al. (2011).

While *L. alpinus* v. *mitis* appears to be quite distantly related to *L. alpinus* v. *alpinus*, it is very closely related to the European *L. lilacinus* (Lasch) Fr., the recently described Mexican *L. cuspidaurantiacus* Montoya, Bandala & Garay-Serr., and to a lesser extent *L. lepidotus* Hesler & A.H. Sm., all of which are also associated with *Alnus* (Figure 2A). *Lactarius lilacinus* is typically more pink to violet in color and has a fruity smell. *Lactarius cuspidaurantiacus* appears to have more yellow-orange basidiomes, more orange colored, distant lamellae, and a *Pelargonium*-like odor (Montoya et al. 2014). *Lactarius lepidotus* has a more gray-brown to beige pileus. Neither *L. lilacinus*, *L. cuspidaurantiacus* or *L. lepidotus* have been reported from the Rocky Mountains.

2. *Lactarius nanus* J. Favre

Figure 4

The following morphological description includes data from Colorado specimens listed in Barge et al. (2016).

Description. Pileus 10–50 mm in diameter, shallowly convex to plano-convex becoming plane to infundibuliform, \pm papillate, smooth, slightly viscid to dry, with whitish glaucous coating when immature, easily rubbing away or fading in age, deep brown to gray-brown to liver brown with lighter and darker areas present, often lighter toward margin, becoming lighter overall upon desiccation; margin straight when young becoming upturned and wavy to convoluted in age. Lamellae adnate to subdecurrent, subdistant, cream when immature becoming dingy cream to light tan in age, sometimes with a faint pinkish tinge, discoloring to dingy tan where damaged. Stipe 5–30 \times 3–15 mm, equal to clavate, central, smooth, dry, at first covered with whitish glaucous coating as in pileus, pale apricot to dingy cream-tan, hollow. Context dingy cream to brownish. Latex scarce to undetectable, watery, white, unchanging. Odor mild. Taste mild to slightly acrid.

Basidiospores 7–10.5(–12) \times 5–8 μ m, $Q = 1.1$ –1.6, subglobose to ellipsoid; ornamentation forming an incomplete reticulum. Pleuromacrocystidia 53–96.5(–114) \times 6.5–11.5 μ m, scattered to abundant, lanceolate; apex acute to moniliform. Cheilo-



Figure 4. *Lactarius nanus*. Collection EB106-13 near *Salix arctica* (pictured), *S. reticulata*, and *S. planifolia*, Beartooth Plateau, Montana, USA. Scale bar: 2 cm. Photo by E. Barge.

macrocystidia $26\text{--}91.5 \times 7.5\text{--}10 \mu\text{m}$, scattered to abundant, subulate to linear; apex acute to rounded.

Ecology and distribution. Widespread in arctic-alpine areas in the Northern Hemisphere with *Salix*. In the GYE and elsewhere in the Rocky Mountains it occurs above tree line in alpine areas with *Salix arctica*, *S. reticulata*, and *S. planifolia*, late summer.

Specimens examined. U.S.A. MONTANA: Carbon County, Beartooth Plateau, Highline Trail, among shrubby and dwarf *Salix*, 7 Aug 1998, CLC1221 (MONT); among *Salix reticulata* and other *Salix* spp., 8 Aug 1998, ZT6422 (MONT); among *Salix arctica*, *S. planifolia* and *S. reticulata*, 18 Aug 2013, EB106-13 (MONT). WYOMING: Park County, Beartooth Plateau, Frozen Lakes, among *Salix arctica*, 21 Aug 1999, CLC1403 (MONT); Park County, Beartooth Plateau, Solifluction Terraces, among *Salix arctica*, *S. planifolia* and *S. reticulata*, 11 Aug 2012, EB125 (MONT).

Discussion. *L. nanus* is morphologically and phylogenetically close to *L. hysginoides* Korhonen & T. Ulvinen (Figure 2A), however, the latter has a more cinnamon-toned pileus, smaller basidiospores ($6.5\text{--}8.5 \times 4.9\text{--}7.1 \mu\text{m}$), and occurs in boreal areas with *Betula*, *Picea* and *Salix*; it has not been reported from the Rocky Mountains. *Lactarius pallidomarginatus* Barge and C.L. Cripps is also similar, however, it is a violet staining species. Differences between *L. nanus* and *L. glyciosmus* (Fr.) Fr. are discussed in the comments section under the latter.

3. *Lactarius glyciosmus* (Fr.) Fr.

Figure 5

The following morphological description includes data from Colorado specimens listed in Barge et al. (2016).

Description. Pileus 15–50 mm in diameter, broadly convex, later becoming plane with or without a depressed center and small papilla, smooth, dry, azonate to lightly zoned especially near the margin, pale gray-brown to violet-brown with a whitish glaucous coating when immature, fading in age; margin incurved when young and remaining so or becoming straight in age. Lamellae adnate to subdecurrent, subdistant to crowded, cream to pale pinkish to pale yellow-orange. Stipe 10–40 × 3–12 mm, equal to slightly clavate, central to eccentric, smooth, dry, buff or pale salmon with a faint glaucous coating at first as in pileus, stuffed, becoming hollow, often white-mycelioid toward the base. Context pale cream. Latex scarce to undetectable, watery, white, unchanging. Odor of coconut. Taste mild to slightly acrid.

Basidiospores 7–9 × 5–7 µm, Q = 1.2–1.4, broadly ellipsoid to ellipsoid, ornamentation forming an incomplete to nearly complete reticulum. Pleuromacrocytidia 45.5–63.5 × 6.5–7.5 µm, scarce to scattered, subclavate to lanceolate; apex rounded to mucronate. Cheilomacrocytidia 33–66 × 5–9 µm, scattered to abundant, cylindrical to subclavate; apex rounded.

Ecology and distribution. Widespread in the Northern Hemisphere in temperate, boreal, and arctic-alpine areas with *Betula*. In the GYE and elsewhere in the Rocky Mountains, it occurs in subalpine and alpine areas with *Betula glandulosa* and possibly other *Betula* spp., late summer.

Specimens examined. U.S.A. MONTANA: Carbon County, Beartooth Plateau, Birch Site, near *Betula glandulosa*, 29 July 1997, CLC1134 = ZT6096 (MONT); 8 Aug 1998, CLC1217 (MONT); 10 Aug 1999, TWO269 (MONT); 19 Aug 1999, CLC1380 (MONT); 13 Aug 2007, ZT12723 (MONT); Carbon County, Hellroaring Plateau, Hellroaring Creek, near *Betula glandulosa*, 9 Aug 2015, EB111-15 (MONT). WYOMING: Park County, Beartooth Plateau, Island Lake, near *Betula glandulosa*, 6 Sept 2015, EB160-15 (MONT).

Discussion. *Lactarius glyciosmus* is very closely related to *L. mammosus* Fr. (Figure 2A), however, the latter has a browner pileus, stouter stature, and ecology with Pinaceae (although it has been reported with *Betula*) (Heilmann-Clausen et al. 1998). *Lactarius vietus* (Fr.) Fr. has a distinct pale zone near the stipe apex, latex that dries greenish gray, longer pleuromacrocytidia (50–90 × 7–12 µm), a mild to slightly fruity odor, and a strongly acrid taste (Heilmann-Clausen et al. 1998). The presence of *L. mammosus* and *L. vietus* in the Rocky Mountains is unclear. *Lactarius trivialis* (Fr.) Fr., which has been reported from the GYE (McKnight 1982), but could not be confirmed, is typically larger with slightly larger basidiospores (7.3–10 × 5.9–7.8 µm), more numerous pleuromacrocytidia, and a mild to acidic-fruity odor (Heilmann-Clausen et al. 1998).



Figure 5. *Lactarius glyciosmus*. Collection EB111-15 under *Betula glandulosa*, Hellroaring Plateau, Montana, USA. Scale bar: 2 cm. Photo by E. Barge.

Lactarius nanus has larger basidiospores ($7\text{--}10.5\text{--}(12) \times 5\text{--}8 \mu\text{m}$), larger and differently shaped pleuromacrocystidia ($53\text{--}96.5\text{--}(114) \times 6.5\text{--}11.5 \mu\text{m}$), a mild odor, and associates with *Salix*.

4. *Lactarius repraesentaneus* Britzelm.

Figure 6

The following morphological description includes data from Colorado specimens listed in Barge et al. (2016).

Description. Pileus 60–100 mm in diameter, convex to plano-convex usually with a depressed center and sometimes with a papilla, smooth at center, becoming increasingly hairy to bearded toward margin, viscid to dry, azonate, surface orange-brown at center, becoming pale yellow-brown to cream toward margin; hairs pale yellow-brown to red-brown; margin strongly incurved, remaining so or becoming nearly straight in age. Lamellae adnate to decurrent, crowded, cream to pale-yellow, staining violet where damaged. Stipe 30–70 \times 15–35 mm, stout, equal to clavate, viscid to dry, cream to pale-yellow, often with numerous dingy yellow, golden yellow, yellow-brown, or light orange-brown scrobicules, hollow. Context white, staining violet where damaged. Latex scarce to abundant, white, becoming violet. Odor spicy-floral, resinous. Taste resinous to slightly acrid.



Figure 6. *Lactarius repraesentaneus*. Collection EB107-13 near krummholz *Picea engelmannii* and *Salix glauca*, Beartooth Plateau, Montana, USA. Scale bar: 2 cm. Photo by E. Barge.

Basidiospores $8\text{--}10.5 \times 6\text{--}8.5 \mu\text{m}$, $Q = 1.1\text{--}1.4$, broadly ellipsoid to ellipsoid; ornamentation forming an incomplete to dense reticulum. Pleuromacrocyttidia $78.5\text{--}145 \times 9\text{--}13 \mu\text{m}$, scattered to abundant, strongly projecting, subfusiform to lanceolate; apex acute to moniliform. Cheilomacrocyttidia $56\text{--}140 \times 7.5\text{--}13 \mu\text{m}$, scattered to abundant, strongly projecting, subfusiform to lanceolate; apex acute moniliform.

Ecology and distribution. Widespread in the Northern Hemisphere in temperate, boreal, and arctic-alpine areas with *Picea*, *Betula*, and possibly *Salix*. In the GYE, it occurs in the spruce-fir and krummholz zone, typically near *Salix glauca* and *Picea engelmannii*, sometimes also intermixed with *Arctostaphylos uva-ursi* and *Betula glandulosa*, late summer.

Specimens examined. U.S.A. MONTANA: Carbon County, Beartooth Plateau, Birch Site, among *Salix glauca* and krummholz *Picea engelmannii*, 12 Aug 2002, CLC1971 (MONT); 13 Aug 2007, CLC2318 (MONT); 18 Aug 2013, EB107-13 (MONT); Carbon County, Hellroaring Plateau, Hellroaring Creek, among *Arctostaphylos uva-ursi*, *Salix glauca* and krummholz *Picea engelmannii*, 9 Aug 2015, EB116-15 (MONT). WYOMING: Park County, Beartooth Plateau, Frozen Lakes, among *Salix glauca* and krummholz *Picea engelmannii*, 21 Aug 1999, CLC1394 (MONT).

Discussion. Due to its conspicuously bearded margin, and violet staining tissue, *L. repraesentaneus* is difficult to confuse with anything else in the Rocky Mountains. It is very closely related to the arctic-alpine *Dryas* and *Salix* associate *L. dryadophilus* Kühner (Figure 2A), however, the latter features a cream to yellow, smoother pileus with hairs restricted to near the margin, and a basidiospore ornamentation consisting of sparse, thinner ridges (Heilmann-Clausen et al. 1998). *Lactarius dryadophilus* has

not been reported from the Rocky Mountains. *Lactarius repraesentaneus* and other violet-staining species such as *L. uvidus* (Fr.) Fr., *L. aspideus* (Fr.) Fr., *L. salicis-reticulatae* Kühner, *L. montanus* (Hesler & A.H. Sm.) Montoya & Bandala, *L. pallidomarginatus*, and *L. brunneoviolaceus* M.P. Christ., etc. occur together in a weakly supported clade, that corresponds with section *Uvidi* (Konr.) Bon (Figure 2A).

5. *Lactarius salicis-reticulatae* Kühner

Figure 7

The following morphological description is based on Barge et al. (2016).

Description. Pileus 20–40 mm in diameter, convex to broadly convex, usually with a depressed center, smooth, viscid to dry, azonate, cream, pale yellow, or pale ocher, sometimes darker toward the center, staining violet where damaged; margin incurved, remaining so or becoming merely downturned in age. Lamellae adnate to subdecurrent, subdistant to distant, cream, pale yellow, or pale orange-ocher often with a pinkish-buff tint, staining violet where damaged. Stipe 15–20 × 10–15 mm, equal to clavate, viscid to dry, cream to pale-yellow, staining violet where damaged, hollow. Context white, staining violet where damaged. Latex scarce to undetectable, watery, white, becoming violet. Odor mild to slightly sweet. Taste mild.

Basidiospores (7–)8.5–11.5 × (7–)8–10 µm, Q = (1–)1.1–1.4, subglobose to ellipsoid; ornamentation forming an incomplete reticulum. Pleuromacrocytidia 76–101.5 × 7.5–11.5 µm, scarce, strongly projecting, subfusiform to fusiform; apex rounded to acute to moniliform. Cheilomacrocytidia 68.5–91.5 × 7.5–10.5 µm, scattered to abundant, strongly projecting, subfusiform to fusiform; apex acute to rounded to moniliform.

Ecology and distribution. Widespread in arctic-alpine areas in the Northern Hemisphere with *Salix*. In the GYE, it occurs in alpine areas with *S. arctica*, *S. reticulata*, as well as shrubby *Salix* spp., sometimes also intermixed with *Dryas octopetala*, late summer.

Specimens examined. U.S.A. MONTANA: Carbon County, Beartooth Plateau, Birch Site, among *Salix reticulata*, 17 Aug 2011, CLC2776 (MONT); Carbon County, Beartooth Plateau, Highline Trail, among dwarf *Salix* spp., 8 Aug 1998, CLC1211 (MONT); Carbon County, Hellroaring Plateau, Hellroaring Creek, under shrubby *Salix* sp., 8 Aug 2015, EB112-15 (MONT); among shrubby *Salix* sp. and *S. reticulata*, 8 Aug 2015, EB113-15 (MONT); among *Dryas octopetala*, shrubby *Salix* sp. and *S. reticulata*, 8 Aug 2015, EB117-15 (MONT), EB118-15 (MONT), EB119-15 (MONT); Sweet Grass County, Crazy Mountains, above Blue and Granite Lakes, among *Salix arctica*, 1 Aug 2015, EB101-15 (MONT). WYOMING: Park County, Beartooth Plateau, Gardner Lake, under shrubby *Salix* sp., 16 Aug 2014, EB0057-14 (MONT).

Discussion. *Lactarius aspideoides* Burl., described from eastern North America is closely related (Figure 2A), however, it is generally larger, with a somewhat zonate pileus. It also has more crowded lamellae that lack salmon-colored hues, slightly smaller



Figure 7. *Lactarius salicis-reticulatae*. Collection EB0057-14 under shrubby *Salix* sp., Beartooth Plateau, Wyoming, USA. Scale bar: 2 cm. Photo by E. Barge.

basidiospores ($7\text{--}10 \times 7\text{--}8\text{--}\mu\text{m}$) with broader ridges, a bitter to slightly acrid taste, and a subalpine ecology (Hesler and Smith 1979). The status of *L. aspideoides* in western North America is unclear. *Lactarius aspideus*, another similar species which was described from northern Europe with *Salix*, has smaller basidiospores ($6.7\text{--}9.5 \times 5.6\text{--}7.8\text{ }\mu\text{m}$) with denser reticulation, more crowded, creamier colored lamellae, and a habitat mainly in subalpine areas (Heilmann-Clausen et al. 1998). The presence of *L. aspideus* in North America is unclear. *Lactarius salicis-herbaceae* Kühner, a closely related (Figure 2A) arctic-alpine *Salix* associate, which has not been reported from the Rocky Mountains, has darker yellow to yellow-brown sporocarps, pale cream to grayish buff lamellae, and more densely reticulate basidiospores (Heilmann-Clausen et al. 1998).

6. *Lactarius pallidomarginatus* Barge & C.L. Cripps

Figure 8

The following morphological description is from Colorado material which includes the holotype and all are listed in Barge et al. (2016)

Description. Pileus 20–50 mm in diameter, convex to broadly convex to plane with or without a depressed center, smooth, subviscid to dry, azonate, blotchy light tan to light brown, developing violet stains, lighter (to cream) toward margin; margin in-



Figure 8. *Lactarius pallidomarginatus*. Collection EB0041 under *Salix planifolia*, San Juan Mountains, Colorado, USA. Scale bar: 2 cm. Photo by E. Barge.

curved when young, remaining so or becoming nearly straight in age. Lamellae adnate to subdecurrent, subdistant to slightly crowded, white to pale yellow-cream, staining violet where damaged. Stipe 10–40 × 5–10 mm, equal to slightly clavate, smooth, dry, white to cream, staining violet where damaged, hollow. Context white to cream, staining violet where damaged. Latex scarce to undetectable, watery, white, staining tissue violet. Odor mild. Taste mild.

Basidiospores 8–10 × 6.5–8 µm, $Q = 1.1–1.4$, broadly ellipsoid to ellipsoid; ornamentation forming an incomplete to dense reticulum. Pleuromacrocytidia 81.5–112 × 9–10 µm, scarce to scattered, cylindrical to lanceolate; apex acute to moniliform. Cheilomacrocytidia 48–101.5 × 7.5–13 µm, scattered, cylindrical to lanceolate; apex acute to moniliform.

Ecology and distribution. Known from only a few alpine localities in the central and southern Rocky Mountains with *Salix planifolia* and possibly also *S. glauca*, late summer.

Specimens examined. U.S.A. WYOMING: Sublette County, Wind River Range, Union Peak, near *Salix glauca*, 22 Aug 1994, ZT5229 (MONT).

Discussion. This species was recently described (Barge et al. 2016) from the alpine zone in Southern Colorado with *Salix*. It is tentatively included here because a collection morphologically matching the type was made by Dr Egon Horak from the Wind

River Range of Wyoming (technically in the southern GYE) with *Salix glauca*. Interestingly, out of all of the species examined thus far, *L. pallidomarginatus* appears most closely related to the bright yellow, violet-staining *Salix* associate *L. aspideus* (Figure 2A), to which it bears little resemblance.

In the Rocky Mountains, *L. pallidomarginatus* is most easily confused with *L. nanus*, *L. glyciosmus*, *L. montanus*, and *L. aff. brunneoviolaceus*. The basidiomes of *L. nanus* do not stain violet where damaged and it produces basidiospores with thicker, more jagged ridges, and macrocystidia with more rounded apices (see Barge et al. 2016). *Lactarius glyciosmus* also does not stain violet where damaged, it has an odor of coconut, slightly smaller basidiospores ($7\text{--}9 \times 5\text{--}7 \mu\text{m}$), and smaller cheilomacrocystidia ($33\text{--}66 \times 5\text{--}9 \mu\text{m}$). The closely related violet-staining *L. montanus* produces basidiomes which are typically much more robust and stain green on the stipe and pileus surface with KOH. *Lactarius montanus* also has a strongly resinous odor and taste, and is mainly subalpine with conifers. See comments under *L. aff. brunneoviolaceus* for features differentiating *L. pallidomarginatus* from that taxon. See Barge et al. (2016) for features differentiating *L. pallidomarginatus* from the European taxon *L. pseudouvidus* Kühner.

7. *Lactarius montanus* (Hesler & A.H. Sm.) Montoya & Bandala

Figure 9

Description. Pileus 30–100 mm in diameter, convex to \pm depressed–convex to infundibuliform, viscid, smooth, azonate to zonate, zones consisting of darker spots arranged concentrically, gray–brown to vinaceous–gray to violet–brown; margin straight to slightly incurved when young, becoming straight and \pm wavy. KOH on pileus cuticle and stipe green. Lamellae adnate to subdecurrent, crowded to subdistant, cream, discoloring violet where damaged with rust brown stains also often present on older material. Stipe 30–70 \times 10–30 mm, equal to clavate, smooth with light tomentum often present especially toward base, dry, white to light brown, discoloring violet and eventually rust brown where damaged, solid, becoming hollow. Context white, discoloring violet and eventually rust brown where damaged. Latex scarce to copious, white, staining tissue violet. Odor strongly resinous. Taste strongly resinous.

Basidiospores $7\text{--}10.5 \times 6\text{--}8 \mu\text{m}$, $Q = 1.2\text{--}1.5$, broadly ellipsoid to ellipsoid; ornamentation forming a broken to partial reticulum. Pleuromacrocystidia $50\text{--}90 \times 5\text{--}10 \mu\text{m}$ scarce to scattered, fusoid–acuminate; apex acute to moniliform. Cheilomacrocystidia $30\text{--}55 \times 4\text{--}9 \mu\text{m}$, scarce, fusoid–acuminate; apex acute to moniliform.

Ecology and distribution. Under conifers in western North America. In the GYE, it occurs in wet areas, often along streams in the montane spruce–fir and krummholz zones, possibly always in the presence of *Picea engelmannii*, summer and fall.

Specimens examined. U.S.A. MONTANA: Madison County, Madison Range, Mirror Lake, swampy area under *Abies lasiocarpa* and *Picea engelmannii*. 3 Sept 2013, EB120-13 (MONT); Madison County, Tobacco Root Mountains, Branham Lakes, under *Picea engelmannii* and *Salix planifolia*, 31 Aug 2013, CLC3001 (MONT), under



Figure 9. *Lactarius montanus*. Top collection EB0072-14 and bottom collection EB0073-14 under *Picea engelmannii*, Tobacco Root Mountains, Montana, USA. Scale bars: 2 cm. Photos by E. Barge.

Picea engelmannii, 1 Sept 2014, EB0071-14 (MONT), EB0072-14 (MONT), EB0073-14 (MONT).

Discussion. In western North America, similar species with violet-staining flesh include *L. pallescens* Hesler & A.H. Sm., *L. californiensis* Hesler & A.H. Sm., and *L.*

cordovaensis Hesler & A.H. Sm. *Lactarius pallescens* and *L. californiensis* are generally more white overall in color, have a slightly acrid to acrid taste (not strongly resinous), and do not stain green in KOH. *Lactarius cordovaensis* has orange–tan lamellae when young, and possibly slightly smaller basidiospores ($7.5\text{--}9.5 \times 6.5\text{--}8 \mu\text{m}$). *Lactarius uvidus*, a European species whose presence has not been confirmed molecularly for North America, typically has a lighter pileus, a bitter taste (not strongly resinous), and does not stain green in KOH. Previous reports of *L. uvidus* from the GYE (McKnight 1982) and the Central and Southern Rocky Mountains in general likely refer to *L. montanus*. See comments under *L. aff. brunneoviolaceus* and *L. pallidomarginatus* for features differentiating *L. montanus* from those taxa.

8. *Lactarius aff. brunneoviolaceus* M.P. Christ.

Figure 10

Description. Pileus 30–60 mm in diameter, convex to slightly depressed–convex, smooth, viscid, faintly zonate, brownish with lavender hues, marbled, whitish near margin; margin incurved and faintly tomentose when young. KOH on pileus cuticle green. Lamellae adnate, crowded, cream, staining violet where damaged. Stipe 25–30 \times 20–25 mm, clavate, smooth, white with hoary coating, staining violet where damaged, hollow. Context not observed. Latex white, unchanging, staining tissue violet. Odor sweet. Taste mild, becoming slightly bitter.

Basidiospores $8\text{--}12 \times 6.5\text{--}8 \mu\text{m}$, $Q = 1.2\text{--}1.5$, broadly ellipsoid to ellipsoid; ornamentation forming an incomplete reticulum. Pleuromacrocyttidia $40\text{--}90 \times 8\text{--}12 \mu\text{m}$, scattered to numerous, subfusiform to fusiform; apex mucronate. Cheilomacrocyttidia $40\text{--}88 \times 7\text{--}12 \mu\text{m}$, numerous, fusiform; apex mucronate.

Ecology and distribution. *Lactarius brunneoviolaceus* occurs in arctic-alpine areas with *Salix*. The species described here was found growing near tree line (krummholz spruce present) on the Beartooth Plateau amongst *Salix reticulata* and *S. planifolia*, late summer.

Specimens examined. U.S.A. MONTANA: Custer County, Beartooth Plateau, Birch Site, among krummholz *Picea engelmannii*, *Salix reticulata* and *Salix planifolia*, 15 Aug 2014, CLC3098 (MONT).

Discussion. More collections and molecular data from this species are needed, however, this is the first report of a species from the GYE, and to the best of our knowledge, North America, bearing affinities to the arctic-alpine *Salix* associate *L. brunneoviolaceus*. Unfortunately, high quality sequence data was not obtained for this collection, and only a small portion of ITS1 could be used for phylogenetic analyses, which placed it with *L. brunneoviolaceus*. Morphologically, it is also a close match, except that the pileus of the species described here stains green with KOH, which is not historically mentioned for *L. brunneoviolaceus*. It is possible that the pileus of *L. brunneoviolaceus* stains green in KOH as well but that it has not been thoroughly tested. In CLC2133 from Norway, another specimen closely matching *L. brunneoviolaceus* which was examined, the pileus (of dried material) also stained green in KOH.



Figure 10. *Lactarius* aff. *brunneoviolaceus*. Collection CLC3098 near *Salix reticulata* (pictured), *S. planifolia*, and krummholz *Picea engelmannii*, Beartooth Plateau, Montana, USA. Scale bar: 2 cm. Photo by C. Cripps.

This species is very closely related to *L. montanus* (Figure 2A), however, the latter features slightly smaller basidiospores ($7\text{--}10.5 \times 6\text{--}8 \mu\text{m}$, although sometimes to $12 \mu\text{m}$ long), and is typically below tree line with conifers. *Lactarius pallidomarginatus*, which is also closely related, has lighter colored, less violaceous pilei that do not stain green in KOH, and smaller basidiospores ($8\text{--}10 \times 6.5\text{--}8 \mu\text{m}$).

9. *Lactarius caespitosus* Hesler & A.H. Sm.

Figure 11

Description. Pileus $35\text{--}100 \mu\text{m}$ in diameter, convex to broadly convex to nearly plane, \pm centrally depressed, viscid to dry, smooth, more or less azonate, cream to pale gray-brown to pale vinaceous-brown, \pm discoloring ochraceous in places; margin incurved when young, becoming straight in age. Lamellae adnate to subdecurrent, crowded to subdistant, white at first, becoming pale creamy buff, \pm discoloring ochraceous to brown where damaged. Stipe $30\text{--}80 \times 10\text{--}35 \text{ mm}$, equal to clavate, viscid to dry, smooth, white, developing ochraceous to faintly violet areas where damaged, solid, becoming hollow. Context white. Latex scarce to undetectable, white, unchanging. Odor mild. Taste acrid.



Figure 11. *Lactarius caespitosus*. Top collection EB0074-14 under *Abies lasiocarpa*, Tobacco Root Mountains, Montana, USA. Bottom collection EB095-15 under *Abies lasiocarpa* and *Picea engelmannii*, Crazy Mountains, Montana, USA. Scale bars: 2 cm. Photos by E. Barge.

Basidiospores $8\text{--}12 \times 7\text{--}9 \mu\text{m}$, $Q = 1.2\text{--}1.4$, broadly ellipsoid to ellipsoid; ornamentation forming a broken to partial reticulum. Pleuromacrocystidia $50\text{--}110 \times 8\text{--}12 \mu\text{m}$, scattered to numerous, subcylindric to fusoid; apex rounded to acute.

Cheilomacrocytidia 40–60 × 8–12 µm, scattered to abundant, clavate to mucronate; apex rounded to acute.

Ecology and distribution. In mountainous areas in western North America with conifers. Hesler and Smith (1979) report it as the common montane *Lactarius* of wet places in the spruce-fir zone of the Rocky Mountains during the summer and early fall. In the GYE, it occurs in wet depressions and near seeps as well as in drier upland sites in the montane spruce-fir zone, possibly always in the presence of *Abies lasiocarpa*, summer to fall.

Specimens examined. U.S.A. MONTANA: Gallatin County, Gallatin Range, East Fork Hyalite Creek, under *Abies lasiocarpa* and *Picea engelmannii*, 25 Aug 2015, EB158-15 (MONT); Gallatin County, Gallatin Range, Windy Pass Trail, under *Abies lasiocarpa* and *Pinus contorta*, 10 Aug 2013, EB102-13 (MONT); Madison County, Madison Range, Big Sky, under *Abies lasiocarpa*, *Picea engelmannii* and *Pinus albicaulis*, 4 Sept 2004, CLC2052 (MONT); Madison County, Madison Range, Mirror Lake, under *Abies lasiocarpa*, 3 Sept 2013, EB121-13 (MONT); Madison County, Tobacco Root Mountains, Branham Lakes, under *Abies lasiocarpa*, 1 Sept 2014, EB0074-14 (MONT); Sweet Grass County, Crazy Mountains, Big Timber Creek, under *Abies lasiocarpa* and *Picea engelmannii*, 1 Aug 2015, EB095-15 (MONT). WYOMING: Park County, Beartooth Plateau, Beartooth Lake Campground, under *Abies lasiocarpa* and *Picea engelmannii*, 5 Sept 2015, EB159-15 (MONT).

Discussion. The European *Abies* associate *L. albocarneus* Britzelm. appears to be closely related (Figure 2B), however, it has a lighter cream to dingy whitish pileus, and slowly yellowing latex and flesh (Heilmann-Clausen et al. 1998). The western North American *L. kauffmanii* Hesler & A.H. Sm. and *L. pseudomucidus* Hesler & A.H. Sm. also appear to be related (Figure 2B). *Lactarius kauffmanii* has a dark to blackish brown pileus (although it can be lighter), smaller basidiospores (7–10 × 6.5–8 µm), and a vinaceous to tan stipe. *Lactarius pseudomucidus* typically has a darker and more slimy pileus, a gray to gray-brown stipe, and smaller basidiospores (7–9 × 6–7 µm). Both *L. kauffmanii* and *L. pseudomucidus* may be confined to Douglas fir/western hemlock forests of the Pacific Northwest (Hesler and Smith 1979). The western North American *L. circellatus* v. *borealis* Hesler & A.H. Sm. is also similar, however, its pileus is pruinose at first and generally darker and faintly zonate to zonate, it has a dry stipe (never viscid as in *L. caespitosus*), and its lamellae are unstaining when cut (Hesler and Smith 1979, Methven 1997). *Lactarius trivialis* is also similar, however it appears to be fairly distantly related (Figure 2A); it has purplish gray to vinaceous colors of the pileus at least when young and smaller basidiospores (7.3–10 × 5.9–7.8 µm) (Heilmann-Clausen et al. 1998). *Lactarius argillaceifolius* Hesler & A.H. Sm. and its varieties are also similar, however, all feature a more purplish gray pileus at least when young (similar to *L. trivialis*) and smaller basidiospores. Variety *argillaceifolius* appears to be confined to oak forests in central and eastern North America, v. *megacarpus* Hesler & A.H. Sm. to oak forests in coastal western North America, and v. *dissimilis* Hesler & A.H. Sm. is represented by only one collection from South Carolina (Hesler and Smith 1979).

10. *Lactarius controversus* Pers.

Figure 12

Description. Pileus 50–120 mm in diameter, depressed–convex to infundibuliform, smooth, viscid to dry, faintly zonate, zones narrow and often more conspicuous near the margin, cream to pale tan–cream, often with tan to rust brown blotches; margin incurved and faintly short–tomentose when young, becoming glabrous and straight to undulating in age. Lamellae adnate to subdecurrent, crowded, creamy pink to pink–tan, discoloring tawny light brown where damaged. Stipe 25–50 × 17–28 mm, short, equal to tapering toward the base, subviscid to dry, smooth, ± scrobiculate, white, discoloring pale yellow to tawny light brown where damaged, solid, becoming hollow. Context white, discoloring tawny light brown. Latex scarce, white, unchanging. Odor faint to slightly spermatic. Taste slowly acrid.

Basidiospores $5.5\text{--}8 \times 4.5\text{--}5.5 \mu\text{m}$, $Q = 1.2\text{--}1.5$, broadly ellipsoid to ellipsoid; ornamentation forming a nearly complete reticulum. Pleuromacrocystidia $25\text{--}50 \times 3\text{--}7 \mu\text{m}$, scarce, cylindric to fusiform; apex acute to moniliform. Cheilomacrocystidia $25\text{--}40 \times 3\text{--}6 \mu\text{m}$, scarce, cylindric to fusiform; apex rounded to acute.

Ecology and distribution. Widely distributed in the northern hemisphere with *Populus* (aspen), *Salix*, and possibly *Betula*. In the GYE it occurs in montane areas near *Populus tremuloides*, summer and fall.

Specimens examined. U.S.A. IDAHO: Teton County, Teton Mountains, under *Populus tremuloides*, CLC290 (MONT). MONTANA: Carbon County, near Red



Figure 12. *Lactarius controversus*. Collection EB110-15 under *Populus tremuloides*, Beartooth Mountains, Montana, USA. Scale bar: 2 cm. Photo by E. Barge.

Lodge, along Rock Creek, under *Populus tremuloides*, 8 Aug 2015, EB110-15 (MONT); Park County, Cinnabar Basin, under *Populus tremuloides*, 18 July 1990, CLC106 (MONT), 2 Aug 1990, CLC116 (MONT), 11 Sept 1991, CLC262 (MONT), 13 July 1992, CLC462 (MONT); Silverbow County, Butte, East Ridge, under *Populus tremuloides*, 15 Aug 1993, CLC699 (MONT). WYOMING: Park County, along Beartooth Highway, under *Populus tremuloides*, 16 Aug 2014, EB0061-14 (MONT).

Discussion. *L. controversus* is fairly easily recognized by its small basidiospores, whitish pileus, pinkish lamellae, acrid taste, and association with aspen. No other species is similar in the Rocky Mountains. While morphological differences between European and GYE material were not detected, there is some molecular divergence, which could indicate separate species, although more material needs to be examined (Figure 2B).

11. *Lactarius olympianus* Hesler & A.H. Sm.

Figure 13

Description. Pileus 20–100 mm in diameter, depressed–convex to infundibuliform, smooth, viscid, zonate, color alternating between rich and dull orange; margin incurved when young becoming straight to wavy in age. Lamellae adnate, crowded to subdistant, cream to dingy yellow, staining orange–brown in age or where damaged.



Figure 13. *Lactarius olympianus*. Collection EB0070-14 under *Picea engelmannii*, Tobacco Root Mountains, Montana, USA. Scale bar: 2 cm. Photo by E. Barge.

Stipe 20–50 × 10–25 mm, equal to tapering toward the base, smooth, white, at first with a white bloom, staining dingy orange–brown where damaged, solid, becoming hollow. Context white. Latex white, unchanging, staining lamellae orange–brown. Odor mild. Taste acrid.

Basidiospores 8–11.5 × 7.5–9.5 µm, $Q = 1.1–1.3$, broadly ellipsoid; ornamentation forming a broken to nearly complete reticulum. Pleuromacrocytidia 20–50 × 3–6 µm, scattered to numerous, fusiform; apex acute to moniliform. Cheilomacrocytidia 20–35 × 3–6 µm, scattered, fusiform; apex acute to moniliform.

Ecology and distribution. In montane conifer forests in western North America. In the GYE, it occurs in seeps and along streams in the spruce–fir zone possibly always in the presence of *Picea engelmannii*.

Specimens examined. U.S.A. MONTANA: Gallatin County, Gallatin Range, Windy Pass Trail, under *Picea engelmannii*, 4 Aug 2012, EB0114 (MONT); Madison County, Tobacco Root Mountains, Branham Lakes, under *Picea engelmannii*, 1 Sept 2014, EB0070-14 (MONT).

Discussion. This species is very close to and likely conspecific with *L. zonarioides* Kühner & Romagn. (Figure 2B), which occurs in similar habitats and was described from Europe (Hesler and Smith 1979).

12. *Lactarius pseudodelicatus* A.H. Sm.

Figure 14

Description. Pileus 70–140 mm in diameter, broadly depressed–convex to broadly infundibuliform, viscid to dry, smooth to matted–tomentose especially near the margin, zonate to faintly zonate, yellow–buff to buff to pale orange–brown, lighter toward the margin; margin incurved and matted–tomentose to cottony–tomentose when young, remaining so or becoming straight and ± glabrous in age. Lamellae subdecurrent to decurrent, many of them branching, crowded, white to pale cream when young, becoming orange–buff to brown to gray–brown in age. Stipe 20–90 × 15–20 mm, tapering toward the base, smooth, white to pale buff, discoloring brown, often with scattered small scrobicules, hollow. Context firm. Latex copious, white, unchanging. Odor pleasant, fruity, of orange–citrus. Taste acrid.

Basidiospores 6–8 × 5–6.5 µm, $Q = 1.2–1.5$, broadly ellipsoid to ellipsoid; ornamentation forming a broken to partial reticulum. Pleuromacrocytidia 60–100 × 7–12 µm, numerous, strongly projecting, fusiform; apex acute. Cheilomacrocytidia 40–61 × 7–10 µm, numerous, strongly projecting, fusiform; apex acute.

Ecology and distribution. Originally described from Michigan under aspen. There are very few subsequent reports. In the GYE, it is here reported with *Populus tremuloides* from the Teton Range of Idaho, summer.

Specimens examined. U.S.A. IDAHO: Teton County, Teton, under *Populus tremuloides*, 27 July 1991, CLC233 (MONT), 24 Aug 1991, CLC296 (MONT), 26 July 1992, CLC512 (MONT).



Figure 14. *Lactarius pseudodelicatus*. Collection CLC512 under *Populus tremuloides*, Teton Range, Idaho, USA. Scale bar: 2 cm. Photo by C. Cripps.

Discussion. This represents the first report of this species from the GYE under the name *L. pseudodelicatus*. It was previously reported from the GYE as *L. cf. zonarius* Fr. in Cripps and Miller (1993) and Cripps (2001). The European *L. zonarius* has slightly larger basidiospores ($6.3\text{--}9.2 \times 5.1\text{--}7.4 \mu\text{m}$), shorter pleuromacrocyttidia ($40\text{--}50 \times 4\text{--}6 \mu\text{m}$), and an ecology with *Quercus* (Heilmann-Clausen et al. 1998).

Molecularly, *L. pseudodelicatus* is very close to a specimen identified as *L. aff. wenquanensis* Y. Wang & Z.X. Xie, from a humid montane rainforest with *Quercus*, *Castanopsis*, and *Lithocarpus echinops* Hjelmq. in Thailand (Figure 2B) (Le et al. 2007). *Lactarius aff. wenquanensis* appears to be less robust with shorter pleuromacrocyttidia ($54\text{--}76 \times 9.5\text{--}12 \mu\text{m}$); it also appears to have a different ecology. *Lactarius wenquanensis* Y. Wang & Z.X. Xie, originally described from China has larger basidiospores ($7.5\text{--}9.5 \times (6\text{--})6.5\text{--}7.5 \mu\text{m}$) and the type was collected under *Picea* and *Abies*, as opposed to aspen (Wang 2007). *Lactarius torminosus* (Schaeff.) Gray is closely related to *L. pseudodelicatus* (Fig. 2B), however, *L. torminosus* typically has smaller basidiomes, a more pink to brick pileus, slightly larger basidiospores ($7.5\text{--}9.6 \times 5.6\text{--}6.9 \mu\text{m}$), and associates with *Betula* (Heilmann-Clausen et al. 1998). *Lactarius pubescens* Fr. is also closely related to *L. pseudodelicatus* (Figure 2B), and both associate with aspen in similar habitats. *Lactarius pubescens* is typically less robust and features a more pale pileus, a non-scrobiculate stipe, and smaller pleuromacrocyttidia ($30\text{--}60 \times 6.5\text{--}10 \mu\text{m}$) (Heilmann-Clausen et al. 1998). *Lactarius delicatus* Burl., described from a deciduous forest in Eastern North America produces white latex that turns yellow (Hesler and Smith 1979).

13. *Lactarius pubescens* Fr.

Figure 15

Description. Pileus 20–100 mm in diameter, depressed–convex to infundibuliform, dry, smooth, hairy toward margin, pale cream to cream or pale pinkish buff; margin bearded, incurved when young becoming straight to wavy in age. Lamellae adnate to decurrent, crowded, pale cream to pale pinkish buff. Stipe 25–50 × 10–15 mm, equal, smooth, dry, pale pinkish buff to pale orange, becoming hollow. Context whitish to pale pinkish buff. Latex not abundant, white, unchanging. Odor fragrant. Taste acrid.

Basidiospores $6\text{--}8.5 \times 4\text{--}6 \mu\text{m}$, $Q = 1.2\text{--}1.5$, broadly ellipsoid to ellipsoid; ornamentation forming an incomplete reticulum. Pleuromacrocystidia $25\text{--}55 \times 7\text{--}10 \mu\text{m}$, scattered, clavate to fusiform; apex acute to moniliform. Cheilomacrocystidia $25\text{--}48 \times 5\text{--}9 \mu\text{m}$, numerous, clavate to fusiform; apex acute to moniliform.

Ecology and distribution. Originally described from Europe and apparently widespread with *Betula* and aspen. It also appears to associate with certain herbaceous plants in alpine areas (China) based on publicly available sequences on GenBank isolated from ectomycorrhizal root tips. In the GYE, it occurs with *Populus tremuloides* and *Betula*.

Specimens examined. U.S.A. MONTANA: Deerlodge County, Anaconda Superfund site, under *Betula* sp., 24 Sept 1996, CLC1045 (MONT); Park County, Gallatin Range, Cinnabar Basin, under *Populus tremuloides*, 1992, CLC539 (MONT); Silver Bow County, 23 Sept 2015, EB300-15 (MONT).



Figure 15. *Lactarius pubescens*. Collection EB303-15 under *Betula* sp., Boise, Idaho, USA. Scale bar: 2 cm. Photo by E. Barge.

Discussion. Morphologically and molecularly (at least at the ITS region), material from the GYE is identical to European material (Figure 2B). In North America, Hesler and Smith (1979) distinguish an *L. pubescens* v. *betulae* (A.H. Sm.) Hesler & A.H. Sm. based on yellowing latex among other subtle features, however yellowing latex was not noted in our collections. Further investigation of *L. pubescens* in North America is warranted. The *Betula* associate *L. torminosus* is similar, and closely related (Figure 2B) but its basidiomes are typically more pink colored and it has larger basidiospores ($7.5\text{--}9.6 \times 5.6\text{--}6.9 \mu\text{m}$) (Heilmann-Clausen et al. 1998). *Lactarius resimus* (Fr.) Fr. is also similar, however it has strongly yellowing latex and flesh and larger basidiospores ($7.0\text{--}9.7 \times 5.3\text{--}7.2 \mu\text{m}$) (Heilmann-Clausen et al. 1998). See comments under *L. pseudodelicatus* for distinguishing features for that taxon.

14. *Lactarius zonarius* v. *riparius* Hesler & A.H. Sm.

Figure 16

Description. Pileus 50–80 mm in diameter, depressed–convex to broadly infundibuliform, viscid to dry, smooth to \pm matted–tomentose toward the margin, zonate to faintly zonate, yellow–brown to orange–brown, \pm lighter toward the margin; margin incurved and matted–tomentose to cottony–tomentose when young, becoming straight and glabrous in age. Lamellae subdecurrent to decurrent, crowded, cream, staining brownish where damaged. Stipe 20–40 \times 15–25 mm, equal to tapering toward the base, smooth,



Figure 16. *Lactarius zonarius* v. *riparius*. Collection CLC2933 under *Populus trichocarpa*, Bozeman, Montana, USA. Scale bar: 2 cm. Photo by E. Barge.

dry, cream, \pm with small pale brown scrobicules. Context very firm, cream. Latex white, unchanging, staining lamellae brownish. Odor sweet, fruity. Taste slowly acrid.

Basidiospores $7\text{--}9 \times 5.5\text{--}7.5 \mu\text{m}$, $Q = 1.2\text{--}1.4$, broadly ellipsoid to ellipsoid; ornamentation forming a highly broken reticulum. Pleuromacrocyttidia $40\text{--}66 \times 4.5\text{--}7.4 \mu\text{m}$, scarce to scattered, narrowly subfusiform to fusiform; apex acute to moniliform. Cheilomacrocyttidia $25\text{--}49 \times 4.5\text{--}5.5 \mu\text{m}$, numerous, narrowly subfusiform to fusiform; apex acute to moniliform.

Ecology and distribution. Originally described and reported from rich wet humus along small streams in hardwood forests in Michigan. There are very few subsequent reports. In the GYE, it occurs in rich, moist riparian areas with *Populus trichocarpa*, summer to fall.

Specimens examined. U.S.A. MONTANA: Gallatin County, Bozeman, along Sourdough Creek, under *Populus trichocarpa*, 2013, CLC2933 (MONT).

Discussion. Molecularly, this taxon is very close to the European *L. evosmus* Kühner & Romagn. (Figure 2B), which also associates with *Populus*, as well as *Quercus*, and occasionally *Salix* (Heilmann-Clausen et al. 1998). It is also close to but distinct from the European *L. zonarius* (Figure 2B). Neither *L. evosmus* or *L. zonarius* feature a bearded pileus margin at any point in sporocarp development. *Lactarius zonarius* v. *riparius* will likely need to be given species level status and a new name, although this hinges on examination of type material.

Similar taxa in western North America include *L. sanmiguelensis* Hesler & A.H. Sm., described from under cottonwood in southwestern Colorado that has a pileus with cinnamon–buff zones on a pinkish buff ground color, the role of felt along the margin is pinkish buff, it does not have a distinctive odor, it is instantly and very strongly acrid, the lamellae are strongly anastomosed, pinkish buff, and do not discolor where injured, and the stipe is longer and pale pinkish buff (Hesler and Smith 1979). Bessette et al. (2009) synonymize *L. sanmiguelensis* and *L. subvillosus* Hesler & A.H. Sm., the latter being undoubtedly different from the species described here. *Lactarius pseudodelicatus* is somewhat similar but it has a longer stipe, smaller basidiospores ($6\text{--}8 \times 5\text{--}6.5 \mu\text{m}$), and much larger pleuromacrocyttidia ($60\text{--}100 \times 7\text{--}12 \mu\text{m}$). *Lactarius psammicola* A.H. Sm. and *L. yazooensis* Hesler & A.H. Sm. are similar species with a more Eastern North American distribution in *Quercus* forests (Hesler and Smith 1979).

15. *Lactarius alnicola* A.H. Sm.

Figure 17

Description. Pileus 60–170 mm in diameter, depressed–convex to infundibuliform, glutinous when wet, with matted fibrils beneath the gluten especially near the margin, \pm faintly zonate, pale creamy yellow to pale yellow–brown to golden yellow–brown, discoloring orange–brown to brown where damaged; margin \pm faintly tomentose, incurved when young, becoming straight to wavy. Lamellae subdecurrent to decurrent, some forked toward the stipe, crowded, pale cream to pale buff, \pm slowly discoloring




Figure 17. *Lactarius alnicola*. Top and bottom collection EB0064-14 under *Picea engelmannii*, Gallatin Range, Montana, USA. Scale bars: 2 cm. Photos by E. Barge.

yellow and eventually orange–brown where damaged. Stipe 20–60 × 30–50 mm, equal to clavate to tapering toward the base, smooth, dry, conspicuously scrobiculate, white, discoloring orange–brown where damaged or in age, solid, becoming hollow. Context firm, white, ± slowly discoloring yellow to orange–brown where damaged. Latex scarce

to undetectable, white, \pm becoming very pale yellow, and slowly staining damaged tissue yellow. Odor mild to sweet. Taste quickly very acrid.

Basidiospores $7.5\text{--}10 \times 6\text{--}8.5 \mu\text{m}$, $Q = 1.2\text{--}1.5$, broadly ellipsoid to ellipsoid; ornamentation forming a partial reticulum. Pleuromacrocyttidia $60\text{--}90 \times 3\text{--}8 \mu\text{m}$, rare, mostly near the pileus and between the lamellae, cylindric to fusiform; apex acute to moniliform. Cheilomacrocyttidia absent.

Ecology and distribution. In western North America and Mexico under conifers. Also reported from California with *Quercus*. In the GYE, *L. alnicola* occurs in wet areas, often along streams in the spruce-fir zone, possibly always in the presence of *Picea engelmannii*, summer and early fall.

Specimens examined. U.S.A. MONTANA: Gallatin County, Gallatin Range, Langhor Road, under *Picea engelmannii*, 27 Aug, EB0064-14 (MONT); Madison County, Tobacco Root Mountains, Branham Lakes, under *Picea engelmannii*, 1 Sept 2014, EB0067-14 (MONT).

Discussion. *Lactarius alnicola* is phylogenetically very closely related to the European *L. scrobiculatus* (Scop.) Fr., and may be conspecific, however more specimens need to be sequenced before making this determination final (Figure 2B). For morphological differences, *L. scrobiculatus* features a heavily bearded pileus margin, and produces white latex which rapidly turns yellow (Kytövuori 1984). *Lactarius alnicola*, and other members of subsection *Scrobiculati* Hesler & A.H. Sm. sensu Kytövuori (1984) form a strongly supported clade in the phylogeny (Figure 2B).

Two varieties of *L. alnicola* have been described: *L. alnicola* v. *pitkinensis* Hesler & A.H. Sm., described from Colorado under aspen and conifers, has cream colored to white basidiomes, an acrid taste, unchanging latex, and unchanging (non-yellowing) flesh; *L. alnicola* v. *pungens* Hesler & A.H. Sm., described from Michigan in mixed forest, has a dull ochraceous to ochraceous–tan, subviscid, glabrous pileus, an acrid taste, and white, unchanging latex, which stains white paper yellow (Hesler and Smith 1979). Neither appear to fit the taxon described here.

16. *Lactarius* aff. *tuomikoskii* Kytöv.

Figure 18

Description. Pileus to 80 mm in diameter, broadly convex with a depressed center, glutinous when wet, with matted fibrils beneath the gluten especially near the margin, azonate, pale yellow–cream to yellow–tan, discoloring brown in places; margin densely bearded, incurved at least when young. Lamellae subdecurrent, crowded, cream to pale yellow–cream, \pm discoloring yellow and eventually rusty brown where damaged. Stipe 30×20 mm, tapering toward the base, smooth, dry, white to pale yellow–tan, with small, faint, dull yellow–tan scrobicules, hollow. Context white, \pm slowly discoloring pale yellow to faintly tan where damaged. Latex scarce, white, becoming very pale yellow, and slowly staining damaged tissue pale yellow. Odor sweet. Taste acrid.




Figure 18. *Lactarius* aff. *tuomikoskii*. Top and bottom collection EB0052-14 under *Picea engelmannii*, Silver Gate, Montana, USA. Scale bars: 2 cm. Photos by E. Barge.

Basidiospores $7\text{--}10 \times 5.5\text{--}7.5 \mu\text{m}$, $Q = 1.2\text{--}1.5$, broadly ellipsoid to ellipsoid; ornamentation forming a broken to nearly complete reticulum. Pleuromacrocytidia $50\text{--}100 \times 7\text{--}12 \mu\text{m}$, abundant, strongly projecting, fusiform to lanceolate; apex acute to moniliform. Cheilomacrocytidia absent.

Ecology and distribution. In montane, rich, moist habitats with *Picea engelmannii*, late summer.

Specimens examined. U.S.A. MONTANA: Park County, Silver Gate, under *Picea engelmannii*, 13 Aug 2014, EB0052-14 (MONT).

Discussion. This species, along with *L. alnicola*, and *L. aff. olivinus* Kytöv. described below, fall in subsection *Scrobiculati* as recognized by Kytövuori (1984). Morphologically, it is quite close to the European *L. tuomikoskii* Kytöv. It is also very close to some interpretations of *L. scrobiculatus* v. *canadensis* A.H. Sm., and *L. gossypinus* Hesler & A.H. Sm., except it has larger basidiospores and pleuromacrocytidia. *Lactarius payettensis* A.H. Sm. is also similar, but differs by having a more robust, heavily scrobiculate stipe and apparently unchanging latex (Hesler and Smith 1979). Both *L. gossypinus* and *L. payettensis* have been previously reported from the GYE (McKnight 1982), however neither were examined for this treatment. Overall, the complex of species surrounding *L. scrobiculatus* in North America, which includes *L. alnicola*, *L. gossypinus*, *L. payettensis*, *L. scrobiculatus* v. *canadensis*, *L. scrobiculatus* v. *montanus* Methven, and the two taxa described here (*L. aff. tuomikoskii*, and *L. aff. olivinus*) appears to be in need of revision, especially in relation to European taxa such as *L. auriolla* Kytöv., *L. leonis* Kytöv., *L. olivinus*, *L. scrobiculatus*, and *L. tuomikoskii* (Kytövuori 1984).

17. *Lactarius aff. olivinus* Kytöv.

Figure 19

Description. Pileus 30–110 mm in diameter, depressed–convex to infundibuliform, viscid when wet, with appressed, agglutinated scales of confluent hairs, especially near the margin, more or less azonate, cream to olive–buff to yellow–tan to orange–tan, \pm darker toward the center; margin incurved and wooly–tomentose when young, often forming a cottony rim, becoming straight and more or less glabrous. Lamellae subdecurrent, crowded, cream, staining yellow where damaged and then fading or eventually becoming pale ochraceous. Stipe 30–45 \times 10–25 mm, equal to clavate to tapering toward the base, smooth, dry, cream to pale yellow–tan to orange–tan, \pm with pale yellow–tan scrobicules, often with a white ring near the apex, solid, becoming hollow. Context white, staining pale yellow at first and then fading. Latex scarce to undetectable, white, becoming yellow, staining tissue yellow and then fading. Odor faintly sweet to faintly spermatic. Taste mild.

Basidiospores $7.5\text{--}10 \times 5.5\text{--}7 \mu\text{m}$, $Q = 1.3\text{--}1.5$, ellipsoid; ornamentation forming a partial reticulum. Pleuromacrocytidia $51\text{--}100 \times 9\text{--}14 \mu\text{m}$, abundant, strongly projecting, fusiform to lanceolate; apex acute to moniliform. Cheilomacrocytidia absent.

Ecology and distribution. In montane, rich, moist habitats with *Picea engelmannii*, mid to late summer.



Figure 19. *Lactarius* aff. *olivinus*. Top collection EB0051-14 and bottom collection EB0050-14 under *Picea engelmannii*, Silver Gate, Montana, USA. Scale bars: 2 cm. Photos by E. Barge.

Specimens examined. U.S.A. MONTANA: Park County, Silver Gate, under *Picea engelmannii*, 13 Aug 2014, EB0050-14 (MONT), EB0051-14 (MONT).

Discussion. This species is morphologically close to the European *L. olivinus* except for the mild rather than acrid taste. Molecularly (Figure 2B), and ecologically, it is also

close, however we refrain from calling it that species until more collections are examined and compared with European material. It is also very close to some interpretations of the North American taxa *L. scrobiculatus* v. *canadensis*, and *L. gossypinus*, except it has larger basidiospores and pleuromacrocytidia. *Lactarius payettensis* is also similar, but it has unchanging latex, and a strongly acrid taste. See comments under *L. aff. tuomikoskii* for additional information.

18. *Lactarius barrowsii* Hesler & A.H. Sm.

Figure 20

Description. Pileus 30–140 mm in diameter, depressed–convex to broadly infundibuliform, viscid to dry, smooth, azonate, cream to dingy pinkish orange to pale yellow–brown to straw yellow, discoloring red and eventually green in age or where damaged; margin incurved with a white bloom when young, becoming straight to upturned and wavy in age. Lamellae adnate to subdecurrent, crowded to subdistant, dingy pinkish orange to creamy yellow–orange, discoloring red and eventually green where damaged. Stipe 15–50 × 10–30 mm, equal to tapering toward the base, dry, smooth, at first with a white bloom, becoming dingy pinkish orange to yellow–brown in places, staining red and eventually green where damaged, solid, becoming hollow. Context whitish, staining red. Latex scarce to undetectable, watery, red, staining flesh red and eventually green. Odor faintly sweet to spermatic in age. Taste mild.

Basidiospores 8.5–10 × 6–7.5 µm, Q = 1.3–1.5, ellipsoid; ornamentation forming a broken reticulum. Pleuromacrocytidia none observed (absent in Hesler and Smith 1979). Cheilomacrocytidia 30–50 × 2–6 µm, scattered, fusoid; apex acute.

Ecology and distribution. In Western North America, previously reported with ponderosa and pinyon pine. In the GYE, it occurs in foothills to low montane dry scrubland, so far reported only from under *Pinus flexilis*, spring and early summer. This is the first report of this species with *Pinus flexilis*.

Specimens examined. U.S.A. MONTANA: Gallatin County, Story Hill, under *Pinus flexilis*, 30 May 2015, EB008-15, EB015-15, EB028-15 (all at MONT); Madison County, Revenue Flats, under *Pinus flexilis*, 1 June 2015, EB037-15 (MONT).

Discussion. This is the first report of *L. barrowsii* from the GYE. *Lactarius barrowsii*, and other members of section *Deliciosi* (Fr.:Fr.) Redeuilh, Verbeken & Walley, such as *L. deliciosus* (L.) Gray, *L. deterrimus* Gröger, and *L. rubrilacteus* Hesler & A.H. Sm. form a monophyletic group (Figure 2B), previously shown by Nuytinck et al. (2007). *Lactarius barrowsii* is morphologically close to the western North American *L. rubrilacteus*. However, in *L. barrowsii*, the lamellae are more orange, the pileus is lighter and more yellow, the margin has a white bloom when young, and the stipe is often whiter. Phylogenetically, *L. barrowsii* is very close to the hypogeous, spring fruiting, pine associated *L. rubriviridis* Desjardin, H.M. Saylor & Thiers, which occurs in similar dry habitats in western North America (Nuytinck et al. 2007).



Figure 20. *Lactarius barrowsii*. Top collection EB008-15 (left and right) and bottom collection EB015-15 under *Pinus flexilis*, Story Hill, Bozeman, Montana, USA. Scale bars: 2 cm. Photos by E. Barge.

19. *Lactarius rubrilacteus* Hesler & A.H. Sm.

Figure 21

Description. Pileus 45–90 mm in diameter, shallowly depressed–convex to broadly infundibuliform, smooth, viscid, zonate, light brown to orange–cream to yellow–orange to orange–brown to carrot to dingy orange, often with green stains; margin incurved when young becoming straight. Lamellae adnate to subdecurrent, crowded to



Figure 21. *Lactarius rubrilacteus*. Collection EB13-040 under *Pseudotsuga menziesii*, Bridger Range, Montana, USA. Scale bar: 2 cm. Photo by E. Barge.

subdistant, cream to pinkish cinnamon to dull pink, discoloring wine red and eventually green where damaged. Stipe 20–40 × 10–25 mm, equal to tapering toward the base, dry, smooth, at first with a light glaucous coating, ± scrobiculate, cream to dull pink to orange–cream, discoloring wine red and eventually green where damaged, solid, becoming hollow. Context white, immediately staining deep wine–red to burgundy. Latex scarce, wine red, staining tissue wine–red to burgundy and eventually green. Odor mild. Taste mild.

Basidiospores 7.5–9.5 × 5–7.5 µm, Q = 1.3–1.5, ellipsoid; ornamentation forming a partial reticulum. Pleuromacrocytidia 40–80 × 4–6 µm, sparse, fusoid; apex acute. Cheilomacrocytidia 30–65 × 6–9 µm, scattered, fusoid; apex acute.

Ecology and distribution. In western North America with *Pseudotsuga menziesii* and *Pinus*. In the GYE this species occurs in foothills to low montane areas, near *Pseudotsuga menziesii*, spring and early summer.

Specimens examined. U.S.A. MONTANA: Gallatin County, Bridger Range, Fairy Lake Road, under *Pseudotsuga menziesii*, 25 June 2013, EB13-040 (MONT); Park County, Absaroka Mountains, George Lake Trailhead, under *Pseudotsuga menziesii*, 27 June 2014, EB900-14 (MONT).

Discussion. The European *L. sanguifluus* (Paulet) Fr. is morphologically similar, however, its lamellae are typically more violet tinged and its stipe is often scrobiculate. Western North American members of the “*L. deliciosus*” group can appear similar, however, their latex does not start out red as in *L. rubrilacteus*. The western North American *Lactarius barrowsii* also has reddish latex, however, it features a whitish to straw colored pileus and ochraceous to pinkish orange lamellae.

20. “*Lactarius deliciosus*” (L.) Gray (group)

Figure 22

Description. Pileus 30–100 mm in diameter, convex to broadly infundibuliform, smooth to faintly areolate, viscid to dry, \pm faintly zonate, pale cream to pale yellow to carrot orange, often with green stains especially in age; margin striate when wet, incurved when young and becoming straight. Lamellae adnate to subdecurrent, crowded to subdistant, creamy orange to carrot orange, discoloring orange to reddish and eventually green where damaged. Stipe 30–70 \times 5–25 mm, equal to clavate to tapering toward the base, viscid to dry, smooth, creamy orange to carrot orange with a white band toward the apex; discoloring carrot orange to dull reddish and eventually green where damaged, hollow. Context yellowish, immediately staining carrot orange to dull reddish and eventually green. Latex scarce, carrot orange, staining tissue carrot orange to dull reddish and eventually green. Odor mild. Taste mild.

Basidiospores 7.5–10.5 \times 5.5–7.5 μm , $Q = 1.3\text{--}1.5$, ellipsoid; ornamentation forming a partial reticulum. Pleuromacrocytidia 40–55 \times 3–6 μm , very sparse, subfusiform; apex moniliform. Cheilomacrocytidia 45–60 \times 4–9 μm , scarce to numerous, subfusiform; apex moniliform.

Ecology and distribution. In North America with conifers. In the GYE, members of this group occur in montane mixed lodgepole/spruce-fir forests, the spruce-fir zone, and the krummholz zone, summer to fall; there are also reports specifically with *Pinus flexilis* (Cripps and Antibus 2011) and *Pinus albicaulis* (Mohatt et al. 2008, Cripps and Antibus 2011), and one collection is reported here with *Arctostaphylos uva-ursi*, well-separated from any conifers.

Specimens examined. U.S.A. MONTANA: Carbon County, Hellroaring Plateau, Hellroaring Creek, among *Arctostaphylos uva-ursi*, 9 Aug 2015, EB107-15 (MONT); Gallatin County, Gallatin Range, East Fork Hyalite Creek, under *Picea engelmannii*, 24 Aug 2014, EB0063-14 (MONT); Madison County, Madison Range, Taylor Creek, under *Abies lasiocarpa*, *Picea engelmannii* and *Pinus contorta*, 25 July 2015, EB089-15 (MONT).

Discussion. While section *Deliciosi* (Fr.:Fr.) Redeuilh, Verbeken & Walley has been resolved as monophyletic (Nuytinck et al. 2007) (Figure 2B), the complex of species surrounding *L. deliciosus* (L.) Gray in North America is poorly resolved. Using molecular techniques, Nuytinck et al. (2007) showed that *L. deliciosus* and *L. deterrimus* Gröger likely do not occur in North America. The collections of the North American varieties of *L. deliciosus* described by Hesler and Smith (1979) that they analyzed formed a clade separate from the European *L. deliciosus* and *L. deterrimus*, but were poorly resolved, with the exception of *L. deliciosus* v. *areolatus* A.H. Smith, which formed a monophyletic group. For descriptions of North American members of this group see Hesler and Smith (1979) and Methven (1997).

Lactarius deliciosus and all of its North American varieties, as well as *L. deterrimus* have been previously reported from the GYE (McKnight 1982, Mohatt et al. 2008,



Figure 22. “*Lactarius deliciosus*” group. Top collection EB107-15 among *Arctostaphylos uva-ursi*, Hell-roaring Plateau, Montana, USA. Bottom collection EB089-15 under *Picea engelmannii*, *Pinus contorta*, and *Abies lasiocarpa*, Madison Range, Montana, USA. Scale bars: 2 cm. Photos by E. Barge.

Cripps 2011, Cripps and Antibus 2011). Specimens from the GYE included in the phylogenetic analysis in the present study fall in the poorly resolved North American “*L. deliciosus*” group including “*L. deliciosus* v. *deterrimus*” (Gröger) Hesler & A.H.

Sm. and *L. deliciosus* v. *olivaceosordidus* Hesler & A.H. Sm. (Figure 2B). Previous reports of *L. thyinos* from the GYE (McKnight 1982), a northeastern North American member of sect. *Deliciosi* that occurs in *Thuja* bogs may refer to the “*L. deliciosus*” group described here. Overall, this group is in need of revision.

21. *Lactarius badiosanguineus* Kühner & Romagn.

Figure 23

Description. Pileus 15–70 mm in diameter, broadly convex to depressed–convex to broadly infundibuliform, \pm umbonate, subviscid to dry, smooth to wrinkled–veined, typically darker when young, liver colored to deep scarlet red to red–brown to orange–brown to dingy orange, conspicuously lighter (to yellow–orange) toward the margin in age; margin \pm striate when wet, slightly incurved to straight when young, remaining straight or becoming slightly wavy in age. Lamellae subdecurrent, crowded to subdistant, cream to creamy yellow to pale tan to pale orange–tan. Stipe 20–70 \times 8–13 mm, equal to slightly clavate, smooth, dry, faintly white–pruinose when young, dingy orange to red–orange, solid, becoming hollow. Context pale tan to red–orange. Latex scarce to abundant, white, unchanging. Odor mild. Taste mild to slightly bitter.

Basidiospores 7–9.5 \times 6–8 μm , $Q = 1.1\text{--}1.4$, subglobose to ellipsoid; ornamentation forming a broken to nearly complete reticulum. Pleuromacrocytidia 40–90 \times 5–10 μm , scattered to numerous, subfusiform; apex obtuse to broadly acute. Cheilomacrocytidia 20–50 \times 5–9 μm , scattered to numerous, subfusiform; apex obtuse to broadly acute.

Ecology and distribution. In Eurasia and North America with *Picea*. In the GYE, this species occurs in moist areas and along streams (although it can also occur on drier upland sites) in the montane spruce–fir zone, possibly always in the presence of *Picea engelmannii*, summer to fall.

Specimens examined. U.S.A. MONTANA: Gallatin County, Gallatin Range, East Fork Hyalite Creek, under *Abies lasiocarpa* and *Picea engelmannii*, 25 Sept 2013, EB200-13 (MONT); under *Abies lasiocarpa*, *Picea engelmannii* and *Pinus contorta*, 24 Aug 2014, EB0062-14 (MONT); Madison County, Tobacco Root Mountains, Branham Lakes, under *Picea engelmannii*; 1 Sept 2014, EB0069-14 (MONT); Sweet Grass County, Crazy Mountains, Big Timber Creek, under *Abies lasiocarpa* and *Picea engelmannii*, 1 Aug 2015, EB094-15 (MONT).

Discussion. This is the first report of this species from the GYE under the name *L. badiosanguineus*. In North American treatments (e.g. Hesler and Smith 1979, Bessette et al. 2009), this species may have gone under the name *L. hepaticus* Plowr. Previous reports of *L. subdulcis* (Pers.) Gray from the GYE (McKnight 1982) likely refer to *L. badiosanguineus*. The European *L. hepaticus* has a duller, browner pileus (although it can be deep red), yellowing latex and flesh, an acrid taste, and it associates with *Pinus* in nutrient-poor, acidic soils (Kränzlin 2005). The European *L. subdulcis* has a more orange–brown to brown pileus (although it can be reddish), a mild to unpleasant



Figure 23. *Lactarius badiosanguineus*. Top collection EB0069-14 under *Picea engelmannii*, Tobacco Root Mountains, Montana, USA. Bottom collection EB200-13 under *Picea engelmannii* and *Abies lasiocarpa*, Gallatin Range, Montana, USA. Scale bars: 2 cm. Photos by E. Barge.

smell, a bitter–acid taste, and an association with *Fagus* (Heilmann-Clausen et al. 1998, Kränzlin 2005). Phylogenetically, *L. badiosanguineus* appears to be fairly closely related to but distinct from both *L. subdulcis* and *L. hepaticus* (Figure 2C).

Lactarius badiosanguineus is closely related to *L. subviscidus* Hesler & A.H. Sm., *L. sphagnetii* (Fr.) Neuhoff, and *L. fulvissimus* Romagn. *Lactarius subviscidus* features typically smaller, more fragile basidiomes with a more uniformly orange pileus, and latex that stains white paper yellow (Hesler and Smith 1979); *L. subviscidus* appears to be confined to the coastal forests of the Pacific Northwest of North America. According to Heilmann-Clausen et al. (1998), the European *L. sphagnetii* features a more distinctly two-toned pileus with a paler margin and darker center (although *L. badiosanguineus* also often displays this character at least in the GYE) as well as more densely reticulate spores with a stronger amyloid reaction. The European *L. fulvissimus* typically features basidiomes which are more orange overall, white latex which often turns yellow, basidiospores with an ornamentation of small pointed warts, and an ecology with deciduous trees such as *Fagus*, *Quercus*, *Tilia*, *Populus*, and *Corylus*, more rarely with conifers (Heilmann-Clausen et al. 1998, Kränzlin 2005).

Lactarius badiosanguineus is also very similar to *L. atrobadius* Hesler & A.H. Sm., reported from the Pacific coastal *Picea sitchensis* belt in North America. *Lactarius atrobadius* appears to have darker basidiomes, however, a more in depth molecular and morphological comparison between the two taxa is warranted. *Lactarius lanceolatus* O.K. Mill. & Laursen is also similar, however it typically produces basidiomes which are overall more orange with larger pleuromacrocytidia ($73.5\text{--}127 \times 6.5\text{--}10\ \mu\text{m}$), and it occurs above tree line with *Salix*. See comments under *L. luculentus* v. *laetus* Hesler & A.H. Sm. for features distinguishing it from *L. badiosanguineus*.

22. *Lactarius rufus* (Scop.) Fr.

Figure 24

Description. Pileus 45–110 mm in diameter, convex to \pm depressed–convex to broadly infundibuliform, smooth, at first covered with a fine bloom becoming \pm faintly areolate in age, subviscid when wet, becoming dry, azonate, deep red–brown to orange–brown, sometimes with gray–brown bands or blotches; margin incurved when young, becoming straight. Lamellae adnate to subdecurrent, subdistant to crowded, cream to dingy orange–tan in age, discoloring dingy brown where damaged. Stipe 40–90 \times 10–15 mm, equal to subclavate, smooth, at first with a fine bloom, dry, cream to tan–pink to red–brown, typically remaining cream toward the base, solid, becoming hollow. Context white to vinaceous–buff. Latex scarce to abundant, white, unchanging. Odor mild. Taste acrid.

Basidiospores $6.8\text{--}9.2 \times 5.1\text{--}7.1\ \mu\text{m}$, $Q = 1.3\text{--}1.5$, ellipsoid; ornamentation forming a complete reticulum. Pleuromacrocytidia $26\text{--}70 \times 6\text{--}12\ \mu\text{m}$, scattered to abundant, subfusiform to fusiform; apex obtuse to mucronate. Cheilomacrocytidia $20\text{--}50 \times 5\text{--}9\ \mu\text{m}$, scattered, subfusiform to fusiform; apex obtuse to broadly acute.

Ecology and distribution. Widely distributed in the northern hemisphere with conifers and *Betula*. In the GYE, it occurs in montane mid-elevation mixed lodgepole/spruce-fir forests up through to high elevation mixed lodgepole/spruce-fir/whitebark pine forests, typically on drier, upland sites, summer to fall.



Figure 24. *Lactarius rufus*. Collection EB0076-14 under *Pinus contorta* and *Picea engelmannii*, Gallatin Range, Montana, USA. Scale bar: 2 cm. Photo by E. Barge.

Specimens examined. U.S.A. MONTANA: Gallatin County, Gallatin Range, Fox Meadow, under *Abies lasiocarpa*, *Picea engelmannii* and *Pinus albicaulis*, 7 Sept 2013, EB125-13 (MONT); Gallatin County, Gallatin Range, Blackmore Trailhead, under *Abies lasiocarpa* and *Pinus contorta*, 15 Sept 2015, EB301-15 (MONT); Gallatin County, Gallatin Range, Lick Creek, under *Picea engelmannii* and *Pinus contorta*, 27 Aug 2014, EB0076-14 (MONT). WYOMING: Park County, Washburn Range, Dunraven Pass, under *Pinus albicaulis* and *Pinus contorta*, 10 Sept 2005, CLC2205 (MONT).

Discussion. *Lactarius rufus* appears to be an extremely morphologically and ecologically variable species, although more research is needed to see if this is backed up molecularly. At the ITS region, specimens from the GYE are molecularly identical to European specimens (*RPB2* data missing for European collections) (Figure 2C). Similar species in the Rocky Mountains include *L. badiosanguineus* and *L. luculentus* v. *laetus*, both of which are less robust and not acrid.

23. *Lactarius luculentus* v. *laetus* Hesler & A.H. Sm.

Figure 25

Description. Pileus 30–50 mm in diameter, broadly convex to broadly infundibuliform, ± umbonate, smooth, viscid when wet but soon dry, yellow–orange to dull orange to brilliant orange to orange–brown; margin straight when young to wavy in age. Lamellae



Figure 25. *Lactarius luculentus* v. *laetus*. Collection EB097-15 under *Abies lasiocarpa* and *Picea engelmannii*, Crazy Mountains, Montana, USA. Scale bar: 2 cm. Photo by E. Barge.

adnate to decurrent, crowded to subdistant, cream to pale yellow–orange. Stipe 20–60 × 5–10 mm, equal to slightly clavate to tapering toward the base, smooth, subviscid to dry, lighter than to concolorous with the pileus, typically getting lighter toward the apex, sometimes with scattered small scrobicules present, solid, becoming hollow. Context cream to pale yellow. Latex scarce, white, unchanging. Odor mild. Taste mild to slightly bitter.

Basidiospores 7–10 × 5.5–8 µm, $Q = 1.2$ –1.5, broadly ellipsoid to ellipsoid; ornamentation forming a very broken reticulum. Pleuromacrocytidia 60–98 × 8–12 µm, scattered to numerous, strongly projecting, subfusiform to fusiform; apex acute. Cheilomacrocytidia none observed, possibly absent.

Ecology and distribution. In western North America with conifers. In the GYE it occurs in the montane spruce–fir zone, summer.

Specimens examined. U.S.A. MONTANA: Sweet Grass County, Crazy Mountains, Big Timber Creek, under *Abies lasiocarpa* and *Picea engelmannii*, 1 Aug 2015, EB097-15 (MONT).

Discussion. This is the first report of this taxon from the GYE under the name *L. luculentus* v. *laetus*. Previous reports of *L. aurantiacus* (Pers.) Gray from subalpine areas in the GYE (McKnight 1982) may refer to *L. luculentus* v. *laetus*.

Variety *laetus* is distinguished from var. *luculentus* by a brighter orange pileus as opposed to orange–brown and a mild to slightly bitter taste as opposed to slightly acrid (Hesler and Smith 1979). At the molecular level, there is some divergence within *L. luculentus* (Figure 2C, see also Barge et al. 2016) and there may be more than one species hiding under the name, although more specimens need to be examined and sequenced before making any nomenclatural changes.

Lactarius luculentus v. *laetus* is very closely related to *L. aurantiacus* and *L. lanceolatus* (Figure 2C) and morphological differences between them are subtle. *Lactarius lanceolatus* appears to have longer and more numerous cheilomacrocytidia ($35.5\text{--}89 \times 5\text{--}10 \mu\text{m}$), longer pleuromacrocytidia ($73.5\text{--}127 \times 6.5\text{--}10 \mu\text{m}$), as well as an alpine ecology with *Salix*. *Lactarius aurantiacus* also appears to have slightly longer pleuromacrocytidia ($68.6\text{--}119.4 \times 7.6\text{--}12.7 \mu\text{m}$), and, while it occupies similar habitats to *L. luculentus* v. *laetus*, it may be absent from western North America outside of Alaska (Barge 2015).

The closely related *Betula* associate *L. lapponicus* Harmaja (syn.: *L. duplicatus* A.H. Sm.) (Figure 2C) has yellowing latex as opposed to white, unchanging latex, and appears to have a more boreal to arctic distribution (Heilmann-Clausen et al. 1998); it has not been reported from the Rocky Mountains. The closely related *L. substriatus* A.H. Sm. and *L. subflammeus* Hesler & A.H. Sm. (Figure 2C) have slightly smaller basidiospores ($6.5\text{--}9 \times 6\text{--}8 \mu\text{m}$, $7.5\text{--}9 \times 6.5\text{--}7.5 \mu\text{m}$ respectively), a slightly acrid taste, and are scarlet when young (Hesler and Smith 1979). The latex of *L. substriatus* also slowly stains pale yellow. Phylogenetic analyses suggest *L. substriatus* and *L. subflammeus* are conspecific, and that yellowing latex is a variable feature within the combined species, although more specimens should be sequenced before making this determination final. Neither of these taxa have been reported from the GYE and their distribution seems to be centered along the Pacific Northwest coast. The morphologically similar *L. subviscidus*, also centered along the Pacific Northwest coast, typically has smaller and more fragile basidiomes, a slightly acrid taste and unchanging latex which stains white paper yellow (Hesler and Smith 1979). In the Rocky Mountains, *L. luculentus* v. *laetus* and *L. badiosanguineus* occur in the same habitat and can easily be confused. However, the latter has slightly smaller basidiospores ($7\text{--}9.5 \times 6\text{--}8 \mu\text{m}$) and typically has a more red-brown pileus which is often darker toward the center and lighter toward the margin.

24. *Lactarius lanceolatus* O.K. Mill. & Laursen

Figure 26

The following description is based on Barge et al. (2016)

Description. Pileus 10–45 mm in diameter, convex to depressed–convex at first, becoming plane to infundibuliform, with or without a small papilla, smooth, sometimes faintly scaly toward the center, viscid to dry, azonate, deep orange–brown to deep orange when immature becoming light to deep orange, often blotchy; margin incurved to straight when immature, becoming straight to upturned and often slightly wavy when mature, \pm slightly crenulate. Lamellae adnate to subdecurrent, slightly crowded, pale cream to pale yellow to pale orange, discoloring brownish orange in age or where damaged. Stipe 10–20 \times 2.5–7.5 mm, equal to clavate, smooth, dry, at first covered by faint whitish pubescence, pale orange, discoloring dingy orange to dingy light brown where damaged, hollow. Context pale orange. Latex scarce to undetectable, watery, white, unchanging. Odor mild. Taste mild.



Figure 26. *Lactarius lanceolatus*. Top collection EB103-15 under shrubby *Salix* spp. and *Betula glandulosa*, Hellroaring Plateau, Montana, USA. Bottom collection EB105-13 among *Salix arctica*, *S. reticulata*, and *S. planifolia*, Beartooth Plateau, Wyoming, USA. Scale bars: 2 cm. Photos by E. Barge.

Basidiospores $8\text{--}10 \times 6\text{--}8 \mu\text{m}$, $Q = 1.1\text{--}1.4$, broadly ellipsoid; ornamentation forming a broken reticulum. Pleuromacrocytidia $73.5\text{--}127 \times 6.5\text{--}10 \mu\text{m}$, scattered to abundant, strongly projecting, fusiform to lanceolate; apex acute to moniliform. Cheilomacrocytidia $35.5\text{--}89 \times 5\text{--}10 \mu\text{m}$, sparse to abundant, strongly projecting, conical to fusiform; apex acute to moniliform.

Ecology and distribution. Widespread in arctic-alpine areas in the northern hemisphere with *Salix*. In the GYE, it occurs above tree line in alpine areas with *Salix reticulata*, *S. arctica*, and *S. planifolia*, late summer.

Specimens examined. U.S.A. MONTANA: Carbon County, Beartooth Plateau, Highline Trail, among dwarf and shrubby *Salix* spp., 1 Aug 1997, ZT6214 (MONT); among *Salix* spp. 7 Aug 1998, ZT6412 (MONT); among *Salix reticulata*, 20 Aug 1999, CLC1389 (MONT); 5 Aug 2008, CLC2358 (MONT); 27 July 1997, CLC1139 (MONT); Carbon County, Hellroaring Plateau, Hellroaring Creek, among *Dryas octopetala* and shrubby *Salix* spp., 8 Aug 2015, EB104-15 (MONT); among *Dryas octopetala*, shrubby *Salix* spp. and *S. reticulata*, 8 Aug 2015, EB105-15 (MONT); among *Betula glandulosa* and shrubby *Salix* spp., 9 Aug 2015, EB103-15 (MONT). WYOMING: Park County, Beartooth Plateau, Solifluction Terraces, among *Salix reticulata*, 20 July 2001, CLC1578 (MONT); 13 Aug 2007, CLC2319 (MONT); among *Salix arctica*, *S. planifolia* and *S. reticulata*, 17 Aug 2013, EB 105-13 (MONT).

Discussion. *Lactarius lanceolatus* is closely related to *L. aurantiacus*, and *L. luculentus*/*L. luculentus* v. *laetus* (Figure 2C). It can be distinguished from *L. luculentus*/*L. luculentus* v. *laetus* in having longer cheilomacrocytidia and an alpine ecology with *Salix*, whereas the latter occur with conifers. Currently, we are unaware of any solid morphological differences between *L. lanceolatus* and *L. aurantiacus*. *Lactarius lapponicus* (syn.: *L. duplicatus*), which occurs in boreal and arctic-alpine areas in the Northern Hemisphere with *Betula*, and possibly other hosts, is also closely related, yet it features strongly yellowing latex (Heilmann-Clausen et al. 1998). Other similar orange taxa in western North America such as *L. substriatus*, *L. subflammeus*, and *L. subviscidus* all occur below tree line with conifers, and their distributions seem to be centered along the Pacific Northwest coast (Hesler and Smith 1979). See comments under *L. badiosanguineus* for features differentiating *L. lanceolatus* from that species.

Discussion

This study reports 20 *Lactarius* species, three varieties, and one unresolved species group from the GYE. *Lactarius alpinus* v. *mitis* with *Alnus incana*, *L. barrowsii* with *Pinus flexilis*, *L. aff. brunneoviolaceus* with *Salix reticulata*, *L. luculentus* v. *laetus* under mixed conifers, *L. aff. olivinus* with *Picea engelmannii*, *L. pseudodelicatus* with *Populus tremuloides*, and *L. aff. tuomikoskii* with *Picea engelmannii* are reported from the GYE for the first time. *Lactarius badiosanguineus* is also reported from the GYE for the first time, and it appears to be fairly common throughout the Rockies with *Picea*, however, it has likely been misidentified in past treatments as *L. hepaticus* (Hesler and Smith 1979, Bessette et al. 2009) and *L. subdulcis* (McKnight 1982).

Several taxonomic problems in need of clarification were encountered. In the Rocky Mountains and North America in general, the group of species surrounding *L. scrobiculatus* (*L. alnicola*, *L. gossypinus*, *L. aff. olivinus*, *L. payettensis*, *L. scrobiculatus* v. *canadensis*,

L. scrobiculatus v. *montanus*, and *L. aff. tuomikoskii*) requires clarification with respect to European taxa such as *L. auriolla*, *L. leonis*, *L. olivinus*, and *L. tuomikoskii* as was already highlighted by Kytövuori (1984). The “*L. deliciosus*” group, whose members are commonly encountered under conifers during the summer and fall in the GYE and elsewhere in western North America is also in need of clarification (Nuytinck et al. 2007). Species delimitation in the *L. aurantiacus*–*L. lanceolatus*–*L. luculentus* complex bares further scrutiny (Barge et al. 2016). The varieties *L. alpinus* v. *mitis*, and *L. zonarius* v. *riparius* may need to be raised to the species level. Potential conspecificity between the North American *L. olympianus* and the European *L. zonarioides* should be investigated. Further, the presence of *L. helvus*, *L. resimus*, *L. torminosus*, and *L. trivialis* in the GYE (McKnight 1982) and North America in general needs clarification (Hesler and Smith 1979).

Barge et al. (2016) reinforced the traditional concept of broad intercontinental distributions for alpine species associated with *Salix* (*L. lanceolatus*, *L. nanus*, *L. salicis-reticulatae*) and *Betula* (*L. glyciosmus*, *L. pubescens*). Based on the results of the present study, broad intercontinental distributions are also suggested for some *Picea* associates (e.g. *L. badiosanguineus*, *L. repraesentaneus*), some aspen associates (*L. controversus*, *L. pubescens*), and *L. rufus*, whose range of host affinities within the GYE remain difficult to untangle. However, it has been shown that at least some regions of the *Lactarius* genome, particularly the ITS region, are relatively slowly evolving (Verbeken and Nuytinck 2013, Barge et al. 2016), thus inclusion of more loci and individuals in phylogenetic analyses would likely reveal finer biogeographic patterns for these taxa. Broad intercontinental distributions have also been confirmed for some western North American arctic-alpine and subalpine members of the ectomycorrhizal genus *Inocybe* (Cripps et al. 2010, Larsson et al. 2014) as well as for some members of the ectomycorrhizal genus *Cortinarius* subg. *Phlegmacium* associated with *P. tremuloides* in western North America (Cripps et al. 2015).

Lactarius species reported from the GYE such as *L. barrowsii* with *Pinus flexilis*, *L. rubrilacteus* with *Pseudotsuga menziesii*, and *L. caespitosus* with *Abies lasiocarpa*, appear to be restricted to western North America. The distribution and host affinities of some species is not clear due in part to taxonomic problems (e.g. “*L. deliciosus*” group) or the relative rarity with which they have been collected and reported (e.g. *L. pallidomarginatus*, *L. pseudodelicatus*, *L. zonarius* v. *riparius*). Interestingly, nearly every major ectomycorrhizal host plant in the GYE has at least one *Lactarius* species associated with it, with some species appearing to be highly host specific (Table 2), highlighting the importance of this genus in the GYE.

Acknowledgments

We thank Vera Evenson, Dr Egon Horak, Jukka Vauras, Dr Ellen Larsson, Dr Pierre-Arthur Moreau, Dr Ursula Peintner, and herbaria G, IB, TU, O, GB, TURA, C, LIP, MICH, DBG, VPI, ZT and NY for loan of specimens. Support from the Montana Institute on Ecosystems graduate enhancement fund is also gratefully acknowledged.

References

- Barge EG (2015) Systematics of *Lactarius* in the Rocky Mountain alpine zone. Master's Thesis, Montana State University, USA.
- Barge EG, Cripps CL, Osmundson TW (2016) Systematics of the ectomycorrhizal genus *Lactarius* in the Rocky Mountains alpine zone. *Mycologia* 108(2): 414–440. doi: 10.3852/15-177
- Bessette AE, Harris DB, Bessette AR (2009) Milk mushrooms of North America: A field identification guide to the genus *Lactarius*. Syracuse University Press, 1–297.
- Buyck B, Hofstetter V, Eberhardt U, Verbeken A, Kauff F (2008) Walking the thin line between *Russula* and *Lactarius*: the dilemma of *Russula* subsect. *Ochricompactae*. *Fungal Diversity* 28: 15–40.
- Buyck B, Hofstetter V, Verbeken A, Walley R (2010) Proposal to conserve *Lactarius* nom. cons. (Basidiomycota) with a conserved type. *Taxon* 59: 295–296.
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* 17: 540–552. doi: 10.1093/oxfordjournals.molbev.a026334
- Cripps CL (2001) Mycorrhizal fungi of aspen forests: natural occurrence and potential applications. In: Shepperd WD, Binkley D, Bartos DL, Stohlgren TJ, Eskew LG (Eds) Proceedings of the symposium on western aspen forests. Rocky Mountain Research Station report RMRS-P-18, USDA Forest Service, Fort Collins, Colorado, USA, 285–298.
- Cripps CL (2003) Native mycorrhizal fungi with aspen on smelter-impacted sites in the Northern Rocky Mountains: occurrence and potential use in reclamation. In: Proceedings of the National Meeting of the American Society of Mining and Reclamation and the 9th Billings Land Reclamation Symposium. Society of Mined Land Reclamation, Lexington, Kentucky, USA, 193–208.
- Cripps CL (2011) Fungi in Yellowstone National Park: a technical report consisting of two volumes. Heritage and Research Center, Yellowstone National Park, USA, 1–114.
- Cripps CL, Antibus RK (2011) Native ectomycorrhizal fungi of limber and whitebark pine: Necessary for forest sustainability? In: Keane RE, Tomback DF, Murray MP, Smith CM (Eds) Proceedings of the high five symposium on the future of high-elevation, five-needle white pines in western North America. Rocky Mountain Research Station report RMRS-P-63, USDA Forest Service, Missoula, Montana, USA, 28–30.
- Cripps CL, Barge EG (2013) Notes on the genus *Lactarius* from the Rocky Mountain alpine zone in regard to Finnish arctic-alpine species. *Karstenia* 53: 29–37.
- Cripps CL, Eddington LH (2005) Distribution of mycorrhizal types among alpine vascular plant families on the Beartooth Plateau, Rocky Mountains, USA, in reference to large-scale patterns in arctic-alpine habitats. *Arctic, Antarctic and Alpine Research* 37: 177–188. doi: 10.1657/1523-0430(2005)037[0177:DOMTAA]2.0.CO;2
- Cripps CL, Eddington LH (2012) What do we know about fungi in Yellowstone National Park? *Yellowstone Science* 20(1): 6–16.
- Cripps CL, Horak E (2008) Checklist and ecology of the Agaricales, Russulales and Boletales in the alpine zone of the Rocky Mountains (Colorado, Montana, Wyoming) at 3000–4000 m A.S.L. *Sommerfeltia* 31: 101–123. doi: 10.2478/v10208-011-0005-5

- Cripps CL, Miller OK Jr (1993) Ectomycorrhizal fungi associated with aspen on three sites in the north-central Rocky Mountains. *Canadian Journal of Botany* 71(11): 1414–1420. doi: 10.1139/b93-170
- Cripps CL, Evenson VS, Kuo M (2016) *The Essential Guide to Rocky Mountain Mushrooms by Habitat*. University of Illinois Press, 1–260.
- Cripps CL, Larsson E, Horak E (2010) Subgenus *Mallocybe* (*Inocybe*) in the Rocky Mountain alpine zone with molecular reference to European arctic-alpine material. *North American Fungi* 5: 97–126.
- Cripps CL, Liimatainen K, Niskanen T, Dima B, Bishop RF, Ammirati JF (2015) Intercontinental distributions of species of *Cortinarius*, subgenus *Phlegmacium*, associated with *Populus* in western North America. *Botany* 93(11): 711–721. doi: 10.1139/cjb-2015-0089
- Geml J, Laursen GA, Timling I, McFarland JM, Booth G, Lennon N, Nusbaum C, Taylor DL (2009) Molecular phylogenetic biodiversity assessment of arctic and boreal ectomycorrhizal *Lactarius* Pers. (Russulales; Basidiomycota) in Alaska, based on soil and sporocarp DNA. *Molecular Ecology* 18: 2213–2227. doi: 10.1111/j.1365-294X.2009.04192.x
- Heilmann-Clausen J, Verbeken A, Vesterholt J (1998) The genus *Lactarius*. *Danish Mycological Society*, 1–287.
- Hesler LR, Smith AH (1960) Studies on *Lactarius*-II. The North American species of sections *Scrobiculus*, *Crocei*, *Theiogali* and *Vellus*. *Brittonia* 12(4): 306–350. doi: 10.2307/2805123
- Hesler LR, Smith AH (1979) *North American species of Lactarius*. University of Michigan Press, 1–841.
- Keiter RB, Boyce MS (1991) *The Greater Yellowstone Ecosystem, redefining America's wilderness heritage*. Yale University Press, 1–442.
- Knight DH (1994) *Mountains and plains: the ecology of Wyoming landscapes*. Yale University Press, 1–352.
- Kränzlin F (2005) *Fungi of Switzerland. Volume 6, Russulaceae*. Verlag Mykologia Luzern, 1–317.
- Kytövuori I (1984) *Lactarius* subsection *Scrobiculati* in NW Europe. *Karstenia* 24: 41–72.
- Larsson E, Vauras J, Cripps CL (2014) *Inocybe leiocephala*, a species with an intercontinental distribution range—disentangling the *I. leiocephala*–*subbrunnea*–*catalaunica* morphological species complex. *Karstenia* 54: 15–39.
- Laursen GA, Ammirati JF (1982) *Lactarii* in Alaskan tundra. In: Laursen GA, Ammirati JF, Redhead SA (Eds) *Arctic and Alpine Mycology 1*. University of Washington Press, 245–281.
- Le HT, Nuytinck J, Verbeken A, Lumyong S, Desjardin DE (2007) *Lactarius* in Northern Thailand: 1. *Lactarius* subgenus *Piperites*. *Fungal Diversity* 24: 173–224.
- Marston RA, Anderson JE (1991) Watersheds and vegetation of the Greater Yellowstone Ecosystem. *Conservation Biology* 5(3): 338–346. doi: 10.1111/j.1523-1739.1991.tb00147.x
- McKnight KH (1982) *Checklist of mushrooms and other fungi of Grand Teton and Yellowstone National Parks*. Grand Teton Library and Museum Association, 1–21.
- Methven AS (1997) *The Agaricales (gilled fungi) of California. 10 Russulaceae II Lactarius*. Mad River Press, 1–79.

- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Proceedings of the Gateway Computing Environments Workshop, New Orleans, Louisiana, USA, 1–8. doi: 10.1109/gce.2010.5676129
- Mohatt KR, Cripps CL, Lavin ML (2008) Ectomycorrhizal fungi of whitebark pine (a tree in peril) revealed by sporocarps and molecular analysis of mycorrhizae from treeline forests in the Greater Yellowstone Ecosystem. *Botany* 86: 14–25. doi: 10.1139/B07-107
- Montoya L, Bandala VM, Garay E (2014) Two new species of *Lactarius* associated with *Alnus acuminata* subsp. *arguta* in Mexico. *Mycologia* 106(5): 949–962. doi: 10.3852/14-006
- Moser MM, McKnight K (1987) Fungi (Agaricales, Russulales) from the alpine zone of Yellowstone National Park and Beartooth Mountains with emphasis on *Cortinarius*. In: Laursen GA, Ammirati JF, Redhead SA (Eds) *Arctic and Alpine Mycology 2*. Plenum Press, 299–318.
- Nuytinck J, Verbeken A, Miller SL (2007) Worldwide phylogeny of *Lactarius* section *Deliciosi* inferred from ITS and glyceraldehyde-3-phosphate dehydrogenase gene sequences. *Mycologia* 99: 820–832. doi: 10.3852/mycologia.99.6.820
- Nuytinck J, Voitek A (2016) *Lactarius alpinus*. *Omphalina* 7(3): 12–14.
- Rochet J, Moreau PA, Manzi S, Gardes M (2011) Comparative phylogenies and host specialization in the alder ectomycorrhizal fungi *Alnicola*, *Alpova* and *Lactarius* (Basidiomycota) in Europe. *BMC Evolutionary Biology* 11(1): 40–50. doi: 10.1186/1471-2148-11-40
- Rosenblad MA, Martín MP, Tedersoo L, Ryberg MK, Larsson E, Wurzbacher C, Abarenkov K, Nilsson RH (2016) Detection of signal recognition particle (SRP) RNAs in the nuclear ribosomal internal transcribed spacer 1 (ITS1) of three lineages of ectomycorrhizal fungi (Agaricomycetes, Basidiomycota). *MycoKeys* 13: 21–33. doi: 10.3897/mycokeys.13.8579
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313. doi: 10.1093/bioinformatics/btu033
- Stöver BC, Müller KF (2010) TreeGraph 2: combining and visualizing evidence from different phylogenetic analyses. *BMC Bioinformatics* 11: 7. doi: 10.1186/1471-2105-11-7
- Stucky B (2012) SeqTrace: a graphical tool for rapidly processing DNA sequencing chromatograms. *Journal of Biomolecular Techniques* 23: 90–93. doi: 10.7171/jbt.12-2303-004
- Verbeken A, Nuytinck J (2013) Not every milkcap is a *Lactarius*. *Scripta Botanica Belgica* 51: 162–168.
- Verbeken A, Stubbe D, van de Putte K, Eberhardt U, Nuytinck J (2014) Tales of the unexpected: angiocarpous representatives of the *Russulaceae* in tropical South East Asia. *Persoonia* 32: 13–24. doi: 10.3767/003158514X679119
- Wang XH (2007) Type studies of *Lactarius* species published from China. *Mycologia* 99(2): 253–268. doi: 10.3852/mycologia.99.2.253
- Wisitrassameewong K, Looney BP, Le HT, De Crop E, Das K, Van de Putte K, Eberhardt U, Jiayu G, Stubbe D, Hyde KD, Verbeken A, Nuytinck J (2016) *Lactarius* subgenus *Russularia* (Basidiomycota, Russulales): novel Asian species, worldwide phylogeny and evolutionary relationships. *Fungal Biology*. doi: 10.1016/j.funbio.2016.08.004